

TITLE OF THE INVENTION

Polymer, Resist Composition and Patterning Process

5 This invention relates to (i) a polymer having a specific structure, (ii) a resist composition comprising the polymer as a base resin, and (iii) a patterning process using the resist composition.

10

BACKGROUND OF THE INVENTION

100
200
300
400
500
600
700
800
900
1000
1100
1200
1300
1400
1500
1600
1700
1800
1900
2000
2100
2200
2300
2400
2500
2600
2700
2800
2900
3000
3100
3200
3300
3400
3500
3600
3700
3800
3900
4000
4100
4200
4300
4400
4500
4600
4700
4800
4900
5000
5100
5200
5300
5400
5500
5600
5700
5800
5900
6000
6100
6200
6300
6400
6500
6600
6700
6800
6900
7000
7100
7200
7300
7400
7500
7600
7700
7800
7900
8000
8100
8200
8300
8400
8500
8600
8700
8800
8900
9000
9100
9200
9300
9400
9500
9600
9700
9800
9900
10000
10100
10200
10300
10400
10500
10600
10700
10800
10900
11000
11100
11200
11300
11400
11500
11600
11700
11800
11900
12000
12100
12200
12300
12400
12500
12600
12700
12800
12900
13000
13100
13200
13300
13400
13500
13600
13700
13800
13900
14000
14100
14200
14300
14400
14500
14600
14700
14800
14900
15000
15100
15200
15300
15400
15500
15600
15700
15800
15900
16000
16100
16200
16300
16400
16500
16600
16700
16800
16900
17000
17100
17200
17300
17400
17500
17600
17700
17800
17900
18000
18100
18200
18300
18400
18500
18600
18700
18800
18900
19000
19100
19200
19300
19400
19500
19600
19700
19800
19900
20000
20100
20200
20300
20400
20500
20600
20700
20800
20900
21000
21100
21200
21300
21400
21500
21600
21700
21800
21900
22000
22100
22200
22300
22400
22500
22600
22700
22800
22900
23000
23100
23200
23300
23400
23500
23600
23700
23800
23900
24000
24100
24200
24300
24400
24500
24600
24700
24800
24900
25000
25100
25200
25300
25400
25500
25600
25700
25800
25900
26000
26100
26200
26300
26400
26500
26600
26700
26800
26900
27000
27100
27200
27300
27400
27500
27600
27700
27800
27900
28000
28100
28200
28300
28400
28500
28600
28700
28800
28900
29000
29100
29200
29300
29400
29500
29600
29700
29800
29900
30000
30100
30200
30300
30400
30500
30600
30700
30800
30900
31000
31100
31200
31300
31400
31500
31600
31700
31800
31900
32000
32100
32200
32300
32400
32500
32600
32700
32800
32900
33000
33100
33200
33300
33400
33500
33600
33700
33800
33900
34000
34100
34200
34300
34400
34500
34600
34700
34800
34900
35000
35100
35200
35300
35400
35500
35600
35700
35800
35900
36000
36100
36200
36300
36400
36500
36600
36700
36800
36900
37000
37100
37200
37300
37400
37500
37600
37700
37800
37900
38000
38100
38200
38300
38400
38500
38600
38700
38800
38900
39000
39100
39200
39300
39400
39500
39600
39700
39800
39900
40000
40100
40200
40300
40400
40500
40600
40700
40800
40900
41000
41100
41200
41300
41400
41500
41600
41700
41800
41900
42000
42100
42200
42300
42400
42500
42600
42700
42800
42900
43000
43100
43200
43300
43400
43500
43600
43700
43800
43900
44000
44100
44200
44300
44400
44500
44600
44700
44800
44900
45000
45100
45200
45300
45400
45500
45600
45700
45800
45900
46000
46100
46200
46300
46400
46500
46600
46700
46800
46900
47000
47100
47200
47300
47400
47500
47600
47700
47800
47900
48000
48100
48200
48300
48400
48500
48600
48700
48800
48900
49000
49100
49200
49300
49400
49500
49600
49700
49800
49900
50000
50100
50200
50300
50400
50500
50600
50700
50800
50900
51000
51100
51200
51300
51400
51500
51600
51700
51800
51900
52000
52100
52200
52300
52400
52500
52600
52700
52800
52900
53000
53100
53200
53300
53400
53500
53600
53700
53800
53900
54000
54100
54200
54300
54400
54500
54600
54700
54800
54900
55000
55100
55200
55300
55400
55500
55600
55700
55800
55900
56000
56100
56200
56300
56400
56500
56600
56700
56800
56900
57000
57100
57200
57300
57400
57500
57600
57700
57800
57900
58000
58100
58200
58300
58400
58500
58600
58700
58800
58900
59000
59100
59200
59300
59400
59500
59600
59700
59800
59900
60000
60100
60200
60300
60400
60500
60600
60700
60800
60900
61000
61100
61200
61300
61400
61500
61600
61700
61800
61900
62000
62100
62200
62300
62400
62500
62600
62700
62800
62900
63000
63100
63200
63300
63400
63500
63600
63700
63800
63900
64000
64100
64200
64300
64400
64500
64600
64700
64800
64900
65000
65100
65200
65300
65400
65500
65600
65700
65800
65900
66000
66100
66200
66300
66400
66500
66600
66700
66800
66900
67000
67100
67200
67300
67400
67500
67600
67700
67800
67900
68000
68100
68200
68300
68400
68500
68600
68700
68800
68900
69000
69100
69200
69300
69400
69500
69600
69700
69800
69900
70000
70100
70200
70300
70400
70500
70600
70700
70800
70900
71000
71100
71200
71300
71400
71500
71600
71700
71800
71900
72000
72100
72200
72300
72400
72500
72600
72700
72800
72900
73000
73100
73200
73300
73400
73500
73600
73700
73800
73900
74000
74100
74200
74300
74400
74500
74600
74700
74800
74900
75000
75100
75200
75300
75400
75500
75600
75700
75800
75900
76000
76100
76200
76300
76400
76500
76600
76700
76800
76900
77000
77100
77200
77300
77400
77500
77600
77700
77800
77900
78000
78100
78200
78300
78400
78500
78600
78700
78800
78900
79000
79100
79200
79300
79400
79500
79600
79700
79800
79900
80000
80100
80200
80300
80400
80500
80600
80700
80800
80900
81000
81100
81200
81300
81400
81500
81600
81700
81800
81900
82000
82100
82200
82300
82400
82500
82600
82700
82800
82900
83000
83100
83200
83300
83400
83500
83600
83700
83800
83900
84000
84100
84200
84300
84400
84500
84600
84700
84800
84900
85000
85100
85200
85300
85400
85500
85600
85700
85800
85900
86000
86100
86200
86300
86400
86500
86600
86700
86800
86900
87000
87100
87200
87300
87400
87500
87600
87700
87800
87900
88000
88100
88200
88300
88400
88500
88600
88700
88800
88900
89000
89100
89200
89300
89400
89500
89600
89700
89800
89900
90000
90100
90200
90300
90400
90500
90600
90700
90800
90900
91000
91100
91200
91300
91400
91500
91600
91700
91800
91900
92000
92100
92200
92300
92400
92500
92600
92700
92800
92900
93000
93100
93200
93300
93400
93500
93600
93700
93800
93900
94000
94100
94200
94300
94400
94500
94600
94700
94800
94900
95000
95100
95200
95300
95400
95500
95600
95700
95800
95900
96000
96100
96200
96300
96400
96500
96600
96700
96800
96900
97000
97100
97200
97300
97400
97500
97600
97700
97800
97900
98000
98100
98200
98300
98400
98500
98600
98700
98800
98900
99000
99100
99200
99300
99400
99500
99600
99700
99800
99900
100000
100100
100200
100300
100400
100500
100600
100700
100800
100900
101000
101100
101200
101300
101400
101500
101600
101700
101800
101900
102000
102100
102200
102300
102400
102500
102600
102700
102800
102900
103000
103100
103200
103300
103400
103500
103600
103700
103800
103900
104000
104100
104200
104300
104400
104500
104600
104700
104800
104900
105000
105100
105200
105300
105400
105500
105600
105700
105800
105900
106000
106100
106200
106300
106400
106500
106600
106700
106800
106900
107000
107100
107200
107300
107400
107500
107600
107700
107800
107900
108000
108100
108200
108300
108400
108500
108600
108700
108800
108900
109000
109100
109200
109300
109400
109500
109600
109700
109800
109900
110000
110100
110200
110300
110400
110500
110600
110700
110800
110900
111000
111100
111200
111300
111400
111500
111600
111700
111800
111900
112000
112100
112200
112300
112400
112500
112600
112700
112800
112900
113000
113100
113200
113300
113400
113500
113600
113700
113800
113900
114000
114100
114200
114300
114400
114500
114600
114700
114800
114900
115000
115100
115200
115300
115400
115500
115600
115700
115800
115900
116000
116100
116200
116300
116400
116500
116600
116700
116800
116900
117000
117100
117200
117300
117400
117500
117600
117700
117800
117900
118000
118100
118200
118300
118400
118500
118600
118700
118800
118900
119000
119100
119200
119300
119400
119500
119600
119700
119800
119900
120000
120100
120200
120300
120400
120500
120600
120700
120800
120900
121000
121100
121200
121300
121400
121500
121600
121700
121800
121900
122000
122100
122200
122300
122400
122500
122600
122700
122800
122900
123000
123100
123200
123300
123400
123500
123600
123700
123800
123900
124000
124100
124200
124300
124400
124500
124600
124700
124800
124900
125000
125100
125200
125300
125400
125500
125600
125700
125800
125900
126000
126100
126200
126300
126400
126500
126600
126700
126800
126900
127000
127100
127200
127300
127400
127500
127600
127700
127800
127900
128000
128100
128200
128300
128400
128500
128600
128700
128800
128900
129000
129100
129200
129300
129400
129500
129600
129700
129800
129900
130000
130100
130200
130300
130400
130500
130600
130700
130800
130900
131000
131100
131200
131300
131400
131500
131600
131700
131800
131900
132000
132100
132200
132300
132400
132500
132600
132700
132800
132900
133000
133100
133200
133300
133400
133500
133600
133700
133800
133900
134000
134100
134200
134300
134400
134500
134600
134700
134800
134900
135000
135100
135200
135300
135400
135500
135600
135700
135800
135900
136000
136100
136200
136300
136400
136500
136600
136700
136800
136900
137000
137100
137200
137300
137400
137500
137600
137700
137800
137900
138000
138100
138200
138300
138400
138500
138600
138700
138800
138900
139000
139100
139200
139300
139400
139500
139600
139700
139800
139900
140000
140100
140200
140300
140400
140500
140600
140700
140800
140900
141000
141100
141200
141300
141400
141500
141600
141700
141800
141900
142000
142100
142200
142300
142400
142500
142600
142700
142800
142900
143000
143100
143200
143300
143400
143500
143600
143700
143800
143900
144000
144100
144200
144300
144400
144500
144600
144700
144800
144900
145000
145100
145200
145300
145400
145500
145600
145700
145800
145900
146000
146100
146200
146300
146400
146500
146600
146700
146800
146900
147000
147100
147200
147300
147400
147500
147600
147700
147800
147900
148000
148100
148200
148300
148400
148500
148600
148700
148800
148900
149000
149100
149200
149300
149400
149500
149600
149700
149800
149900
150000
150100
150200
150300
150400
150500
150600
150700
150800
150900
151000
151100
151200
151300
151400
151500
151600
151700
151800
151900
152000
152100
152200
152300
152400
152500
152600
152700
152800
152900
153000
153100
153200
153300
153400
153500
153600
153700
153800
153900
154000
154100
154200
154300
154400
154500
154600
154700
154800
154900
155000
155100
155200
155300
155400
155500
155600
155700
155800
155900
156000
156100
156200
156300
156400
156500
156600
156700
156800
156900
157000
157100
157200
157300
157400
157500
157600
157700
157800
157900
158000
158100
158200
158300
158400
158500
158600
158700
158800
158900
159000
159100
159200
159300
159400
159500
159600
159700
159800
159900
160000
160100
160200
160300
160400
160500
160600
160700
160800
160900
161000
161100
161200
161300
161400
161500
161600
161700
161800
161900
162000
162100
162200
162300
162400
162500
162600
162700
162800
162900
163000
163100
163200
163300
163400
163500
163600
163700
163800
163900
164000
164100
164200
164300
164400
164500
164600
164700
164800
164

in their backbone have a practically acceptable level of etching resistance because the resin backbone is robust, but are very low in sensitivity and resolution because the reactivity of acid-decomposable protective groups is
5 extremely low as compared with those on the (meth)acrylic polymers. Since the backbone of the resin is too robust, substrate adhesion is poor. These compositions are thus impractical as well.

Both the (meth)acrylic and alicyclic backbone systems
10 commonly have the problem of pattern disruption due to swelling of resist film. Resist compositions based on these systems have been designed so as to improve their resolution performance by increasing the difference in dissolution rate before and after exposure, and as a consequence, they
15 eventually become highly hydrophobic. Highly hydrophobic resist compositions, when applied as a film and processed with a developer, can maintain the film tenaciously in unexposed regions and allow the film to be instantaneously dissolved in over-exposed regions, while relatively broad
20 exposed regions therebetween allow penetration of the developer, but are kept undissolved, that is, swollen. At the very small pattern size for which ArF excimer laser is actually used, those resist compositions which allow adjacent pattern strips to be joined together and disrupted
25 on account of swelling are rejected. While a finer pattern rule is being demanded, there is a need to have a resist material which is not only satisfactory in sensitivity, resolution, and etching resistance, but fully restrained from swelling.

SUMMARY OF THE INVENTION

Therefore, an object of the present invention is to provide (i) a polymer having improved reactivity, robustness and substrate adhesion as well as minimized swell during
35 development, (ii) a resist composition comprising the polymer as a base resin, which has a higher resolution and

carbon atoms which may contain a hetero atom; each of R^5 to R^8 is hydrogen, a hydroxyl group or a straight, branched or cyclic, monovalent hydrocarbon group of 1 to 15 carbon atoms which may contain a hetero atom, at least one of R^5 to R^8 contains a hetero atom, any two of R^5 to R^8 may bond together to form a ring, wherein the ring-forming two R's together represent a straight, branched or cyclic, divalent hydrocarbon group of 1 to 15 carbon atoms which may contain a hetero atom; R^9 and R^{10} each are hydrogen or methyl; each of R^{11} to R^{14} is hydrogen or a straight, branched or cyclic, monovalent hydrocarbon group of 1 to 15 carbon atoms which may contain a hetero atom, a pair of R^{11} and R^{12} , a pair of R^{11} or R^{12} and R^{13} , a pair of R^{11} or R^{12} and R^{14} , or a pair of R^{13} and R^{14} may bond together to form a ring, wherein each pair represents a straight, branched or cyclic, divalent hydrocarbon group of 1 to 15 carbon atoms which may contain a hetero atom; R^{15} is hydrogen, methyl or $CH_2CO_2R^{17}$; $R^{15'}$ is hydrogen, methyl or $CH_2CO_2R^{17'}$; $R^{15''}$ is hydrogen, methyl or $CH_2CO_2R^{17''}$; $R^{15'''}$ is hydrogen, methyl or $CH_2CO_2R^{17'''}$; R^{16} is hydrogen, methyl or CO_2R^{17} ; $R^{16'}$ is hydrogen, methyl or $CO_2R^{17'}$; $R^{16''}$ is hydrogen, methyl or $CO_2R^{17''}$; $R^{16'''}$ is hydrogen, methyl or $CO_2R^{17'''}$; R^{17} , $R^{17'}$, $R^{17''}$ and $R^{17'''}$ may be identical or different between R^{15} and R^{16} , between $R^{15'}$ and $R^{16'}$, between $R^{15''}$ and $R^{16''}$, and between $R^{15'''}$ and $R^{16'''}$, respectively, and each is a straight, branched or cyclic alkyl group of 1 to 15 carbon atoms; R^{18} is hydrogen or a monovalent hydrocarbon group of 1 to 15 carbon atoms containing a carboxyl or hydroxyl group; R^{19} is a monovalent hydrocarbon group of 2 to 15 carbon atoms containing at least one partial structure selected from the group consisting of ether, aldehyde, ketone, ester, carbonate, acid anhydride, amide and imide; R^{20} is a polycyclic hydrocarbon group of 7 to 15 carbon atoms or an alkyl group containing a polycyclic hydrocarbon group; R^{21} is an acid labile group; k is 0 or 1; x1, x2, x3, a, b, c and d represent a molar compositional ratio of the recurring units associated therewith, satisfying $x_1+x_2+x_3+a+b+c+d = 1$, x1, x2, x3, a, b and c are numbers inclusive of 0, d is a number

of more than 0, all of x_1 , x_2 and x_3 are not equal to 0 at the same time. Preferably, the acid labile group represented by R^{21} in formula (1) is a tertiary alkyl group having a cyclic structure.

5 In a second aspect, the invention provides a resist composition comprising the polymer defined above.

In a third aspect, the invention provides a process for forming a resist pattern comprising the steps of applying the resist composition defined above onto a
10 substrate to form a coating; heat treating the coating and then exposing it to high-energy radiation or electron beams through a photo mask; and optionally heat treating the exposed coating and developing it with a developer.

207322102302
The polymer comprising recurring units of formula (1)
15 is a polymer based on a (meth)acrylic acid derivative, which is copolymerized with at least one of a vinyl ether compound, an allyl ether compound, a norbornene derivative containing an oxygen atom within its ring, and a tetracyclododecene derivative containing an oxygen atom
20 within its ring. As pointed out above, polymers of (meth)acrylic acid derivatives have good sensitivity and resolution, but poor etching resistance, partly because of many ester bonds and a low carbon density. By changing some of the copolymerized units to the above-mentioned units, the
25 polymer of the invention has succeeded in improving etching resistance without sacrificing the sensitivity and resolution characteristic of the (meth)acrylic acid derivative polymers. And the inventive polymer drastically suppresses the occurrence of undesired phenomena including
30 developer repellency, swelling upon development, and pattern disruption as often found when alicyclic rings are introduced into the backbone. Therefore, a resist composition using the inventive polymer as a base resin satisfies all the performance factors of sensitivity,
35 resolution and etching resistance, is fully restrained from swelling upon development, and is thus very useful in forming micropatterns.

It is noted that hybrid polymers analogous to the inventive polymer are known in the art, including hybrid polymers in which alicyclic olefin compound (e.g., norbornene)-maleic anhydride alternating copolymers are

5 copolymerized with (meth)acrylic acid derivative units and hybrid polymers in which vinyl ether-maleic anhydride alternating copolymers are copolymerized with (meth)acrylic acid derivative units. It has been proposed to use these hybrid polymers as the base resin in resist compositions.

10 These hybrid polymers contain maleic anhydride as an essential unit. However, the maleic anhydride units can adversely affect etching resistance on account of a high oxygen content, and react with water within the system during shelf storage, with the performance of the resist


15 composition being markedly degraded by this ring opening reaction. The inventive polymer, which is obtained by introducing a vinyl ether compound, an allyl ether compound and an oxygen-containing alicyclic olefin compound into a (meth)acrylic acid derivative polymer, rather than

20 introducing maleic anhydride, is not only improved in sensitivity, resolution and etching resistance, but also has excellent storage stability.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

25 Polymer

Polymers or high molecular weight compounds comprising recurring units of the following general formula (1) according to the invention are novel. The polymers have a weight average molecular weight of 1,000 to 500,000.



Herein R^1 and R^2 each are hydrogen or methyl.

R^3 and R^4 each are hydrogen or a straight, branched or cyclic, monovalent hydrocarbon group of 1 to 15 carbon atoms which may contain a hetero atom. Examples include alkyl groups such as methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, tert-amyl, n-pentyl, n-hexyl, cyclopentyl, cyclohexyl, ethylcyclopentyl, butylcyclopentyl, ethylcyclohexyl, butylcyclohexyl, adamantyl, ethyladamantyl, and butyladamantyl. Also included are the foregoing groups in which some of the hydrogen atoms are replaced by a group of hetero atoms, or a group of hetero atoms intervenes between adjacent carbon atoms, that is, groups having a partial structure such as alcohol, ether, aldehyde, ketone, carboxylic acid, ester, carboxylic anhydride, amide, sulfonic acid, sulfonate or sulfonamide. R^3 and R^4 may bond together to form a ring. In that event, a combination of R^3 with R^4 represents a straight, branched or cyclic, divalent hydrocarbon group of 1 to 15 carbon atoms which may contain a hetero atom, examples of which are divalent groups obtained by eliminating a hydrogen atom at an arbitrary position from the groups exemplified above as the monovalent hydrocarbon groups which may contain a hetero atom.

Each of R^5 to R^8 is hydrogen, a hydroxyl group or a straight, branched or cyclic, monovalent hydrocarbon group of 1 to 15 carbon atoms which may contain a hetero atom, and at least one of R^5 to R^8 contains a hetero atom. Examples of the hydrocarbon group are the same as exemplified above for R^3 and R^4 . Any two of R^5 to R^8 may bond together to form a ring. In that event, the combination of ring-forming two R's represents a straight, branched or cyclic, divalent hydrocarbon group of 1 to 15 carbon atoms which may contain a hetero atom, examples of which are the same as exemplified above for R^3 and R^4 .

R^9 and R^{10} each are hydrogen or methyl.

Each of R^{11} to R^{14} is hydrogen or a straight, branched or cyclic, monovalent hydrocarbon group of 1 to 15 carbon atoms which may contain a hetero atom, examples of which are the same as exemplified above for R^3 and R^4 . A pair of R^{11} and R^{12} , a pair of R^{11} or R^{12} and R^{13} , a pair of R^{11} or R^{12} and R^{14} , or a pair of R^{13} and R^{14} may bond together to form a ring. In that event, each pair represents a straight, branched or cyclic, divalent hydrocarbon group of 1 to 15 carbon atoms which may contain a hetero atom, examples of which are the same as exemplified above for R^3 and R^4 .

R^{15} is hydrogen, methyl or $\text{CH}_2\text{CO}_2R^{17}$, $R^{15'}$ is hydrogen, methyl or $\text{CH}_2\text{CO}_2R^{17'}$, $R^{15''}$ is hydrogen, methyl or $\text{CH}_2\text{CO}_2R^{17''}$, $R^{15'''}$ is hydrogen, methyl or $\text{CH}_2\text{CO}_2R^{17'''}$, R^{16} is hydrogen, methyl or CO_2R^{17} , $R^{16'}$ is hydrogen, methyl or $\text{CO}_2R^{17'}$, $R^{16''}$ is hydrogen, methyl or $\text{CO}_2R^{17''}$, and $R^{16'''}$ is hydrogen, methyl or $\text{CO}_2R^{17'''}$. R^{17} , $R^{17'}$, $R^{17''}$ and $R^{17'''}$ may be identical or different between R^{15} and R^{16} , between $R^{15'}$ and $R^{16'}$, between $R^{15''}$ and $R^{16''}$, and between $R^{15'''}$ and $R^{16'''}$, respectively, and each is a straight, branched or cyclic alkyl group of 1 to 15 carbon atoms, examples of which are the same as exemplified above for R^3 and R^4 .

R^{18} is hydrogen or a monovalent hydrocarbon group of 1 to 15 carbon atoms containing a carboxyl or hydroxyl group. Examples include carboxyethyl, carboxybutyl, carboxycyclopentyl, carboxycyclohexyl, carboxynorbornyl, carboxyadamantyl, hydroxyethyl, hydroxybutyl,

hydroxycyclopentyl, hydroxycyclohexyl, hydroxynorbornyl, and hydroxyadamantyl.

R^{19} is a monovalent hydrocarbon group of 2 to 15 carbon atoms containing at least one partial structure selected from among ether, aldehyde, ketone, ester, carbonate, acid anhydride, amide and imide. Examples include methoxymethyl, methoxyethoxymethyl, 7-oxabicyclo[2.2.1]heptan-2-ylmethyl, 2-oxooxolan-3-yl, 2-oxooxolan-4-yl, 4,4-dimethyl-2-oxooxolan-3-yl, 4-methyl-2-oxooxolan-4-yl, 2-oxo-1,3-dioxolan-4-ylmethyl, and 5-methyl-2-oxooxolan-5-yl.

R^{20} is a polycyclic hydrocarbon group of 7 to 15 carbon atoms or an alkyl group containing a polycyclic hydrocarbon group. Examples include norbornyl, bicyclo[3.3.1]nonyl, tricyclo[5.2.1.0^{2,6}]decyl, adamantyl, ethyladamantyl, butyladamantyl, norbornylmethyl, and adamantylmethyl.


R^{21} is an acid labile group.

The subscript k is equal to 0 or 1. The subscripts x_1 , x_2 , x_3 , a , b , c and d represent a molar compositional ratio of the recurring units associated therewith, satisfying $x_1+x_2+x_3+a+b+c+d = 1$, x_1 , x_2 , x_3 , a , b and c are numbers inclusive of 0, d is a number of more than 0, all of x_1 , x_2 and x_3 are not equal to 0 at the same time. The preferred ranges of x_1 , x_2 , x_3 , a , b , c and d are given below.


| | Preferred range | More preferred range |
|---------|-----------------|----------------------|
| x_1 : | from 0 to 0.7 | from 0 to 0.6 |
| x_2 : | from 0 to 0.7 | from 0 to 0.6 |
| x_3 : | from 0 to 0.7 | from 0 to 0.6 |
| a : | from 0 to 0.4 | from 0 to 0.3 |
| b : | from 0 to 0.5 | from 0 to 0.4 |
| c : | from 0 to 0.4 | from 0 to 0.3 |
| d : | from 0.1 to 0.8 | from 0.2 to 0.7 |

The sum of $x_1+x_2+x_3$ is preferably $0 < x_1+x_2+x_3 \leq 0.9$, more preferably $0.05 \leq x_1+x_2+x_3 \leq 0.7$, and even more preferably $0.1 \leq x_1+x_2+x_3 \leq 0.6$.

The acid labile groups represented by R^{21} may be selected from a variety of such groups. Examples of the acid labile group are groups of the following general formulae (L1) to (L4), tertiary alkyl groups of 4 to 20 carbon atoms, preferably 4 to 15 carbon atoms, trialkylsilyl groups in which each alkyl moiety has 1 to 6 carbon atoms, and oxoalkyl groups of 4 to 20 carbon atoms.



In these formulae and throughout the specification, the broken line denotes a free valence bond. R^{L01} and R^{L02} are hydrogen or straight, branched or cyclic alkyl groups of 1 to 18 carbon atoms, preferably 1 to 10 carbon atoms. Exemplary alkyl groups include methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, cyclopentyl, cyclohexyl, 2-ethylhexyl, and n-octyl. R^{L03} is a monovalent hydrocarbon group of 1 to 18 carbon atoms, preferably 1 to 10 carbon atoms, which may contain a hetero atom such as oxygen, examples of which include unsubstituted straight, branched or cyclic alkyl groups and straight, branched or cyclic alkyl groups in which some hydrogen atoms are replaced by hydroxyl, alkoxy, oxo, amino, alkylamino or the like. Illustrative examples are the substituted alkyl groups shown below.



A pair of R^{L01} and R^{L02} , R^{L01} and R^{L03} , or R^{L02} and R^{L03} may form a ring. Each of R^{L01} , R^{L02} and R^{L03} is a straight or branched alkylene group of 1 to 18 carbon atoms, preferably 1 to 10 carbon atoms when they form a ring.

5 R^{L04} is a tertiary alkyl group of 4 to 20 carbon atoms, preferably 4 to 15 carbon atoms, a trialkylsilyl group in which each alkyl moiety has 1 to 6 carbon atoms, an oxoalkyl group of 4 to 20 carbon atoms, or a group of formula (L1). Exemplary tertiary alkyl groups are tert-butyl, tert-amyl, 10 1,1-diethylpropyl, 2-cyclopentylpropan-2-yl, 2-cyclohexylpropan-2-yl, 2-(bicyclo[2.2.1]heptan-2-yl)propan-2-yl, 2-(adamantan-1-yl)propan-2-yl, 1-ethylcyclopentyl, 1-butylcyclopentyl, 1-ethylcyclohexyl, 1-butylcyclohexyl, 1-ethyl-2-cyclopentenyl, 1-ethyl-2-cyclohexenyl, 2-methyl-2-adamantyl, and 2-ethyl-2-adamantyl. Exemplary trialkylsilyl groups are trimethylsilyl, triethylsilyl, and dimethyl-tert-butylsilyl. Exemplary oxoalkyl groups are 3-oxocyclohexyl, 4-methyl-2-oxooxan-4-yl, and 5-methyl-2-oxooxolan-5-yl. Letter y is an integer of 0 to 6.


15 R^{L05} is a monovalent hydrocarbon group of 1 to 8 carbon atoms which may contain a hetero atom or a substituted or unsubstituted aryl group of 6 to 20 carbon atoms. Examples of the monovalent hydrocarbon group which may contain a hetero atom include straight, branched or cyclic alkyl groups such as methyl, ethyl, propyl, isopropyl, n-butyl, 20 sec-butyl, tert-butyl, tert-amyl, n-pentyl, n-hexyl, cyclopentyl, and cyclohexyl, and substituted groups in which some hydrogen atoms on the foregoing groups are substituted with hydroxyl, alkoxy, carboxy, alkoxycarbonyl, oxo, amino, 25 alkylamino, cyano, mercapto, alkylthio, sulfo or other groups. Exemplary aryl groups are phenyl, methylphenyl, naphthyl, anthryl, phenanthryl, and pyrenyl. Letter m is equal to 0 or 1, n is equal to 0, 1, 2 or 3, and $2m+n$ is equal to 2 or 3.

30 R^{L06} is a monovalent hydrocarbon group of 1 to 8 carbon atoms which may contain a hetero atom or a substituted or

unsubstituted aryl group of 6 to 20 carbon atoms. Examples of these groups are the same as exemplified for R^{L05} .

R^{L07} to R^{L16} independently represent hydrogen or monovalent hydrocarbon groups of 1 to 15 carbon atoms which may contain a hetero atom. Exemplary hydrocarbon groups are straight, branched or cyclic alkyl groups such as methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, tert-amyl, n-pentyl, n-hexyl, n-octyl, n-nonyl, n-decyl, cyclopentyl, cyclohexyl, cyclopentylmethyl, cyclopentylethyl, cyclopentylbutyl, cyclohexylmethyl, cyclohexylethyl and cyclohexylbutyl, and substituted ones of these groups in which some hydrogen atoms are replaced by hydroxyl, alkoxy, carboxy, alkoxycarbonyl, oxo, amino, alkylamino, cyano, mercapto, alkylthio, sulfo or other groups. Alternatively, R^{L07} to R^{L16} , taken together, form a ring (for example, a pair of R^{L07} and R^{L08} , R^{L07} and R^{L09} , R^{L08} and R^{L10} , R^{L09} and R^{L10} , R^{L11} and R^{L12} , R^{L13} and R^{L14} , or a similar pair form a ring). Each of R^{L07} to R^{L16} represents a divalent C_1 - C_{15} hydrocarbon group which may contain a hetero atom, when they form a ring, examples of which are the ones exemplified above for the monovalent hydrocarbon groups, with one hydrogen atom being eliminated. Two of R^{L07} to R^{L16} which are attached to adjoining carbon atoms (for example, a pair of R^{L07} and R^{L09} , R^{L09} and R^{L15} , R^{L13} and R^{L15} , or a similar pair) may bond together directly to form a double bond.

Of the acid labile groups of formula (L1), the straight and branched ones are exemplified by the following groups.




Of the acid labile groups of formula (L1), the cyclic ones are, for example, tetrahydrofuran-2-yl, 2-methyltetrahydrofuran-2-yl, tetrahydropyran-2-yl, and 2-methyltetrahydropyran-2-yl.

5 Examples of the acid labile groups of formula (L2) include tert-butoxycarbonyl, tert-butoxycarbonylmethyl, tert-amylloxycarbonyl, tert-amylloxycarbonylmethyl, 1,1-diethylpropyloxycarbonyl, 1,1-diethylpropyloxycarbonylmethyl, 1-ethylcyclopentyloxycarbonyl, 1-ethylcyclopentyloxy-
10 carbonylmethyl, 1-ethyl-2-cyclopentenylloxycarbonyl, 1-ethyl-2-cyclopentenylloxycarbonylmethyl, 1-ethoxyethoxycarbonylmethyl, 2-tetrahydropyranyloxycarbonylmethyl, and 2-tetrahydrofuranyloxycarbonylmethyl groups.

15 Examples of the acid labile groups of formula (L3) include 1-methylcyclopentyl, 1-ethylcyclopentyl, 1-n-propylcyclopentyl, 1-isopropylcyclopentyl, 1-n-butylcyclopentyl, 1-sec-butylcyclopentyl, 1-methylcyclohexyl, 1-ethylcyclohexyl, 3-methyl-1-cyclopenten-3-yl, 3-ethyl-1-cyclopenten-3-yl, 3-methyl-1-cyclohexen-3-yl, and 3-ethyl-1-cyclohexen-3-yl groups.
20

The acid labile groups of formula (L4) are exemplified by the following groups.




25 Examples of the tertiary alkyl groups of 4 to 20 carbon atoms, trialkylsilyl groups in which each alkyl moiety has 1 to 6 carbon atoms, and oxoalkyl groups of 4 to 20 carbon atoms are as exemplified for R^{L04} .


Of the groups exemplified as the acid labile group of R^{21} , tertiary alkyl groups having a cyclic structure are
30 especially preferred. This is because the relatively low


deblocking reactivity of tertiary alkyl groups under acidic conditions makes the polymer fully stable during storage and because the inclusion of cyclic structure makes the polymer more rigid and hence, well resistant to etching.


- 5 Illustrative, non-limiting examples of the unit to be introduced into the polymer of formula (1) in the molar compositional ratio x1 are shown below.




- 10 Illustrative, non-limiting examples of the unit to be introduced into the polymer of formula (1) in the molar compositional ratio x2 are shown below.









Illustrative, non-limiting examples of the unit to be introduced into the polymer of formula (1) in the molar
 5 compositional ratio x3 are shown below.



Illustrative, non-limiting examples of the unit to be introduced into the polymer of formula (1) in the molar compositional ratio "a" are shown below.




Illustrative, non-limiting examples of the unit to be introduced into the polymer of formula (1) in the molar compositional ratio "b" are shown below.




5

Illustrative, non-limiting examples of the unit to be introduced into the polymer of formula (1) in the molar compositional ratio "c" are shown below.



Illustrative, non-limiting examples of the unit to be introduced into the polymer of formula (1) in the molar compositional ratio "d" are shown below.




5

Each of the recurring units described above is to impart any appropriate properties such as dissolution contrast, developer affinity, substrate adhesion and etching resistance to a resist composition into which the polymer is


formulated. By properly adjusting the content of such recurring units, the performance of the resist composition can be finely adjusted.

The polymers of the invention have a weight average molecular weight of about 1,000 to 500,000, and preferably about 3,000 to 100,000, as measured by gel permeation chromatography (GPC) using a polystyrene standard. Outside the range, the etching resistance may become extremely low and the resolution may become low because a substantial difference in rate of dissolution before and after exposure is lost.


The polymer of the invention can be prepared through copolymerization reaction using one, two or three members selected from compounds of the following general formulae (2) to (4) as first to third monomers, a compound of the following general formula (8) as a next monomer, and optionally, one, two or three members selected from compounds of the following general formulae (5) to (7) as subsequent monomers.




(2)




(3)




(4)




(5)



(6)



(7)



(8)

Herein, R^1 to R^{21} , $R^{15'}$, $R^{15''}$, $R^{16'}$, $R^{16''}$, x_1 , x_2 , x_3 , a , b , c and d are as defined above.

By properly adjusting the proportion of the respective monomers used in the copolymerization reaction, the polymer
5 can be tailored so that it may exert the preferred performance when blended in resist compositions.

In addition to (i) the monomer or monomers of formulas (2) to (4), (ii) the monomer of formula (8), and (iii) the monomer or monomers of formulae (5) to (7), the polymer of
10 the invention may have copolymerized therewith (iv) another monomer having a carbon-to-carbon double bond other than (i) to (iii). Examples of the additional monomer (iv) include substituted acrylic acid esters such as methyl methacrylate, methyl crotonate, dimethyl maleate, and dimethyl itaconate,
15 unsaturated carboxylic acids such as maleic acid, fumaric acid and itaconic acid, and unsaturated acid anhydrides such as itaconic anhydride.

A variety of copolymerization reaction methods may be used for the preparation of the polymer according to the
20 invention. The preferred polymerization reaction is radical polymerization.

For radical polymerization, preferred reaction conditions include (a) a solvent selected from among hydrocarbons such as benzene, ethers such as tetrahydrofuran,
25 alcohols such as ethanol, and ketones such as methyl isobutyl ketone, (b) a polymerization initiator selected from azo compounds such as 2,2'-azobisisobutyronitrile and peroxides such as benzoyl peroxide and lauroyl peroxide, (c) a temperature of about 0°C to about 100°C, and (d) a time of
30 about 1/2 hour to about 48 hours. Reaction conditions outside the described range may be employed if desired.

Resist composition

Since the polymer of the invention is useful as the base resin of a resist composition, the other aspect of the
35 invention provides a resist composition, especially a chemically amplified positive resist composition, comprising the polymer. Typically, the resist composition contains the

polymer, a photoacid generator, and an organic solvent, and other optional components.


Photoacid generator

The photoacid generator is a compound capable of generating an acid upon exposure to high energy radiation or electron beams and includes the following:


- (i) onium salts of the formula (P1a-1), (P1a-2) or (P1b),
- (ii) diazomethane derivatives of the formula (P2),
- (iii) glyoxime derivatives of the formula (P3),
- (iv) bissulfone derivatives of the formula (P4),
- (v) sulfonic acid esters of N-hydroxyimide compounds of the formula (P5),
- (vi) β -ketosulfonic acid derivatives,
- (vii) disulfone derivatives,
- (viii) nitrobenzylsulfonate derivatives, and
- (ix) sulfonate derivatives.

These photoacid generators are described in detail.

(i) Onium salts of formula (P1a-1), (P1a-2) or (P1b):



P1a-1




P1a-2

Herein, R^{101a} , R^{101b} , and R^{101c} independently represent straight, branched or cyclic alkyl, alkenyl, oxoalkyl or oxoalkenyl groups of 1 to 12 carbon atoms, aryl groups of 6 to 20 carbon atoms, or aralkyl or aryloxoalkyl groups of 7 to 12 carbon atoms, wherein some or all of the hydrogen atoms may be replaced by alkoxy or other groups. Also, R^{101b} and R^{101c} , taken together, may form a ring. R^{101b} and R^{101c} each are alkylene groups of 1 to 6 carbon atoms when they form a ring. K^- is a non-nucleophilic counter ion.

R^{101a} , R^{101b} , and R^{101c} may be the same or different and are illustrated below. Exemplary alkyl groups include methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, pentyl, hexyl, heptyl, octyl, cyclopentyl, cyclohexyl,

cycloheptyl, cyclopropylmethyl, 4-methylcyclohexyl, cyclohexylmethyl, norbornyl, and adamantyl. Exemplary alkenyl groups include vinyl, allyl, propenyl, butenyl, hexenyl, and cyclohexenyl. Exemplary oxoalkyl groups include 2-oxocyclopentyl and 2-oxocyclohexyl as well as 2-oxopropyl, 2-cyclopentyl-2-oxoethyl, 2-cyclohexyl-2-oxoethyl, and 2-(4-methylcyclohexyl)-2-oxoethyl. Exemplary aryl groups include phenyl and naphthyl; alkoxyphenyl groups such as p-methoxyphenyl, m-methoxyphenyl, o-methoxyphenyl, ethoxyphenyl, p-tert-butoxyphenyl, and m-tert-butoxyphenyl; alkylphenyl groups such as 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, ethylphenyl, 4-tert-butylphenyl, 4-butylphenyl, and dimethylphenyl; alkyl naphthyl groups such as methyl naphthyl and ethyl naphthyl; alkoxy naphthyl groups such as methoxy naphthyl and ethoxy naphthyl; dialkyl naphthyl groups such as dimethyl naphthyl and diethyl naphthyl; and dialkoxy naphthyl groups such as dimethoxy naphthyl and diethoxy naphthyl. Exemplary aralkyl groups include benzyl, phenylethyl, and phenethyl. Exemplary aryloxoalkyl groups are 2-aryl-2-oxoethyl groups such as 2-phenyl-2-oxoethyl, 2-(1-naphthyl)-2-oxoethyl, and 2-(2-naphthyl)-2-oxoethyl. Examples of the non-nucleophilic counter ion represented by K⁺ include halide ions such as chloride and bromide ions, fluoroalkylsulfonate ions such as triflate, 1,1,1-trifluoroethanesulfonate, and nonafluorobutanesulfonate, arylsulfonate ions such as tosylate, benzenesulfonate, 4-fluorobenzenesulfonate, and 1,2,3,4,5-pentafluorobenzene-sulfonate, and alkylsulfonate ions such as mesylate and butanesulfonate.




P1b

Herein, R^{102a} and R^{102b} independently represent straight, branched or cyclic alkyl groups of 1 to 8 carbon atoms. R¹⁰³ represents a straight, branched or cyclic alkylene groups of

1 to 10 carbon atoms. R^{104a} and R^{104b} independently represent 2-oxoalkyl groups of 3 to 7 carbon atoms. K^+ is a non-nucleophilic counter ion.

Illustrative of the groups represented by R^{102a} and R^{102b} are methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, pentyl, hexyl, heptyl, octyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, 4-methylcyclohexyl, and cyclohexylmethyl. Illustrative of the groups represented by R^{103} are methylene, ethylene, propylene, butylene, pentylene, hexylene, heptylene, octylene, nonylene, 1,4-cyclohexylene, 1,2-cyclohexylene, 1,3-cyclopentylene, 1,4-cyclooctylene, and 1,4-cyclohexanedimethylene. Illustrative of the groups represented by R^{104a} and R^{104b} are 2-oxopropyl, 2-oxocyclopentyl, 2-oxocyclohexyl, and 2-oxocycloheptyl. Illustrative examples of the counter ion represented by K^+ are the same as exemplified for formulae (P1a-1) and (P1a-2). (ii) Diazomethane derivatives of formula (P2)




P2

Herein, R^{105} and R^{106} independently represent straight, branched or cyclic alkyl or halogenated alkyl groups of 1 to 12 carbon atoms, aryl or halogenated aryl groups of 6 to 20 carbon atoms, or aralkyl groups of 7 to 12 carbon atoms.

Of the groups represented by R^{105} and R^{106} , exemplary alkyl groups include methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, pentyl, hexyl, heptyl, octyl, amyl, cyclopentyl, cyclohexyl, cycloheptyl, norbornyl, and adamantyl. Exemplary halogenated alkyl groups include trifluoromethyl, 1,1,1-trifluoroethyl, 1,1,1-trichloroethyl, and nonafluorobutyl. Exemplary aryl groups include phenyl; alkoxyphenyl groups such as p-methoxyphenyl, m-methoxyphenyl, o-methoxyphenyl, ethoxyphenyl, p-tert-butoxyphenyl, and m-tert-butoxyphenyl; and alkylphenyl groups such as 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, ethylphenyl, 4-tert-butylphenyl, 4-butylphenyl, and dimethylphenyl. Exemplary

halogenated aryl groups include fluorophenyl, chlorophenyl, and 1,2,3,4,5-pentafluorophenyl. Exemplary aralkyl groups include benzyl and phenethyl.

(iii) Glyoxime derivatives of formula (P3)



P3

5


Herein, R^{107} , R^{108} , and R^{109} independently represent straight, branched or cyclic alkyl or halogenated alkyl groups of 1 to 12 carbon atoms, aryl or halogenated aryl groups of 6 to 20 carbon atoms, or aralkyl groups of 7 to 12 carbon atoms. Also, R^{108} and R^{109} , taken together, may form a ring. R^{108} and R^{109} each are straight or branched alkylene groups of 1 to 6 carbon atoms when they form a ring.

10

Illustrative examples of the alkyl, halogenated alkyl, aryl, halogenated aryl, and aralkyl groups represented by R^{107} , R^{108} , and R^{109} are the same as exemplified for R^{105} and R^{106} . Examples of the alkylene groups represented by R^{108} and R^{109} include methylene, ethylene, propylene, butylene, and hexylene.

15

(iv) Bissulfone derivatives of formula (P4)




P4

20

Herein, R^{101a} and R^{101b} are as defined above.

(v) Sulfonic acid esters of N-hydroxyimide compounds of formula (P5)



P5

Herein, R^{110} is an arylene group of 6 to 10 carbon atoms, alkylene group of 1 to 6 carbon atoms, or alkenylene group of 2 to 6 carbon atoms wherein some or all of the hydrogen atoms may be replaced by straight or branched alkyl or alkoxy groups of 1 to 4 carbon atoms, nitro, acetyl, or phenyl groups. R^{111} is a straight, branched or cyclic alkyl group of 1 to 8 carbon atoms, alkenyl, alkoxyalkyl, phenyl or naphthyl group wherein some or all of the hydrogen atoms may be replaced by alkyl or alkoxy groups of 1 to 4 carbon atoms, phenyl groups (which may have substituted thereon an alkyl or alkoxy of 1 to 4 carbon atoms, nitro, or acetyl group), hetero-aromatic groups of 3 to 5 carbon atoms, or chlorine or fluorine atoms.

Of the groups represented by R^{110} , exemplary arylene groups include 1,2-phenylene and 1,8-naphthylene; exemplary alkylene groups include methylene, ethylene, trimethylene, tetramethylene, phenylethylene, and norbornane-2,3-diyl; and exemplary alkenylene groups include 1,2-vinylene, 1-phenyl-1,2-vinylene, and 5-norbornene-2,3-diyl. Of the groups represented by R^{111} , exemplary alkyl groups are as exemplified for R^{101a} to R^{101c} ; exemplary alkenyl groups include vinyl, 1-propenyl, allyl, 1-butenyl, 3-butenyl, isoprenyl, 1-pentenyl, 3-pentenyl, 4-pentenyl, dimethylallyl, 1-hexenyl, 3-hexenyl, 5-hexenyl, 1-heptenyl, 3-heptenyl, 6-heptenyl, and 7-octenyl; and exemplary alkoxyalkyl groups include methoxymethyl, ethoxymethyl, propoxymethyl, butoxymethyl, pentyloxymethyl, hexyloxymethyl, heptyloxymethyl, methoxyethyl, ethoxyethyl, propoxyethyl, butoxyethyl, pentyloxyethyl, hexyloxyethyl, methoxypropyl, ethoxypropyl, propoxypropyl, butoxypropyl, methoxybutyl, ethoxybutyl, propoxybutyl, methoxypentyl, ethoxypentyl, methoxyhexyl, and methoxyheptyl.

Of the substituents on these groups, the alkyl groups of 1 to 4 carbon atoms include methyl, ethyl, propyl, isopropyl, n-butyl, isobutyl and tert-butyl; the alkoxy groups of 1 to 4 carbon atoms include methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, isobutoxy, and tert-butoxy;

the phenyl groups which may have substituted thereon an alkyl or alkoxy of 1 to 4 carbon atoms, nitro, or acetyl group include phenyl, tolyl, p-tert-butoxyphenyl, p-acetylphenyl and p-nitrophenyl; the hetero-aromatic groups of 3 to 5 carbon atoms include pyridyl and furyl.

Illustrative examples of the photoacid generator include:

onium salts such as diphenyliodonium trifluoromethanesulfonate, (p-tert-butoxyphenyl)phenyliodonium trifluoromethanesulfonate, diphenyliodonium p-toluenesulfonate, (p-tert-butoxyphenyl)phenyliodonium p-toluenesulfonate, triphenylsulfonium trifluoromethanesulfonate, (p-tert-butoxyphenyl)diphenylsulfonium trifluoromethanesulfonate, bis(p-tert-butoxyphenyl)phenylsulfonium trifluoromethanesulfonate, tris(p-tert-butoxyphenyl)sulfonium trifluoromethanesulfonate, triphenylsulfonium p-toluenesulfonate, (p-tert-butoxyphenyl)diphenylsulfonium p-toluenesulfonate, bis(p-tert-butoxyphenyl)phenylsulfonium p-toluenesulfonate, tris(p-tert-butoxyphenyl)sulfonium p-toluenesulfonate, triphenylsulfonium nonafluorobutanesulfonate, triphenylsulfonium butanesulfonate, trimethylsulfonium trifluoromethanesulfonate, trimethylsulfonium p-toluenesulfonate, cyclohexylmethyl(2-oxocyclohexyl)sulfonium trifluoromethanesulfonate, cyclohexylmethyl(2-oxocyclohexyl)sulfonium p-toluenesulfonate, dimethylphenylsulfonium trifluoromethanesulfonate, dimethylphenylsulfonium p-toluenesulfonate, dicyclohexylphenylsulfonium trifluoromethanesulfonate, dicyclohexylphenylsulfonium p-toluenesulfonate, trinaphthylsulfonium trifluoromethanesulfonate, cyclohexylmethyl(2-oxocyclohexyl)sulfonium trifluoromethanesulfonate, (2-norbornyl)methyl(2-oxocyclohexyl)sulfonium trifluoromethanesulfonate, ethylenebis[methyl(2-oxocyclopentyl)sulfonium trifluoromethanesulfonate], and 1,2'-naphthylcarbonylmethyl-tetrahydrothiophenium triflate;

diazomethane derivatives such as bis(benzenesulfonyl)-diazomethane, bis(p-toluenesulfonyl)diazomethane, bis(xylenesulfonyl)diazomethane, bis(cyclohexylsulfonyl)-

10073223-021302
diazomethane, bis(cyclopentylsulfonyl)diazomethane, bis(n-butylsulfonyl)diazomethane, bis(isobutylsulfonyl)diazomethane, bis(sec-butylsulfonyl)diazomethane, bis(n-propylsulfonyl)diazomethane, bis(isopropylsulfonyl)diazomethane, 5 bis(tert-butylsulfonyl)diazomethane, bis(n-amylsulfonyl)diazomethane, bis(isoamylsulfonyl)diazomethane, bis(sec-amylsulfonyl)diazomethane, bis(tert-amylsulfonyl)diazomethane, 1-cyclohexylsulfonyl-1-(tert-butylsulfonyl)diazomethane, 1-cyclohexylsulfonyl-1-(tert-amylsulfonyl)diazomethane, 10 and 1-tert-amylsulfonyl-1-(tert-butylsulfonyl)diazomethane;

glyoxime derivatives such as bis-O-(p-toluene-sulfonyl)- α -dimethylglyoxime, bis-O-(p-toluenesulfonyl)- α -diphenylglyoxime, bis-O-(p-toluenesulfonyl)- α -dicyclohexylglyoxime, 15 bis-O-(p-toluenesulfonyl)-2,3-pentanedione-glyoxime, bis-O-(p-toluenesulfonyl)-2-methyl-3,4-pentanedione-glyoxime, bis-O-(n-butan-sulfonyl)- α -dimethylglyoxime, bis-O-(n-butan-sulfonyl)- α -diphenylglyoxime, bis-O-(n-butan-sulfonyl)- α -dicyclohexylglyoxime, bis-O-(n-butan-sulfonyl)-2,3-pentanedione-glyoxime, 20 bis-O-(n-butan-sulfonyl)-2-methyl-3,4-pentanedione-glyoxime, bis-O-(methanesulfonyl)- α -dimethylglyoxime, bis-O-(trifluoromethanesulfonyl)- α -dimethylglyoxime, bis-O-(1,1,1-trifluoroethanesulfonyl)- α -dimethylglyoxime, bis-O-(tert-butan-sulfonyl)- α -dimethylglyoxime, 25 bis-O-(perfluorooctanesulfonyl)- α -dimethylglyoxime, bis-O-(cyclohexanesulfonyl)- α -dimethylglyoxime, bis-O-(benzenesulfonyl)- α -dimethylglyoxime, bis-O-(p-fluorobenzenesulfonyl)- α -dimethylglyoxime, bis-O-(p-tert-butylbenzenesulfonyl)- α -dimethylglyoxime, bis-O-(xylenesulfonyl)- α -dimethylglyoxime, 30 and bis-O-(camphorsulfonyl)- α -dimethylglyoxime;

bissulfone derivatives such as bisnaphthylsulfonylmethane, bistrifluoromethylsulfonylmethane, bismethyl-

sulfonylmethane, bisethylsulfonylmethane, bispropylsulfonylmethane, bisisopropylsulfonylmethane, bis-p-toluenesulfonylmethane, and bisbenzenesulfonylmethane;

5 β -ketosulfone derivatives such as 2-cyclohexyl-carbonyl-2-(p-toluenesulfonyl)propane and 2-isopropyl-carbonyl-2-(p-toluenesulfonyl)propane;

disulfone derivatives such as diphenyl disulfone and dicyclohexyl disulfone;

10 nitrobenzyl sulfonate derivatives such as 2,6-dinitrobenzyl p-toluenesulfonate and 2,4-dinitrobenzyl p-toluenesulfonate;

15 sulfonic acid ester derivatives such as 1,2,3-tris-(methanesulfonyloxy)benzene, 1,2,3-tris(trifluoromethanesulfonyloxy)benzene, and 1,2,3-tris(p-toluenesulfonyloxy)benzene; and

20 sulfonic acid esters of N-hydroxyimides such as N-hydroxysuccinimide methanesulfonate, N-hydroxysuccinimide trifluoromethanesulfonate, N-hydroxysuccinimide ethanesulfonate, N-hydroxysuccinimide 1-propanesulfonate, N-hydroxysuccinimide 2-propanesulfonate, N-hydroxysuccinimide 1-pentanesulfonate, N-hydroxysuccinimide 1-octanesulfonate, N-hydroxysuccinimide p-toluenesulfonate, N-hydroxysuccinimide p-methoxybenzenesulfonate, N-hydroxysuccinimide 2-chloroethanesulfonate, N-hydroxysuccinimide benzenesulfonate, 25 N-hydroxysuccinimide 2,4,6-trimethylbenzenesulfonate, N-hydroxysuccinimide 1-naphthalenesulfonate, N-hydroxysuccinimide 2-naphthalenesulfonate, N-hydroxy-2-phenylsuccinimide methanesulfonate, N-hydroxymaleimide methanesulfonate, N-hydroxymaleimide ethanesulfonate, N-hydroxy-2-phenylmaleimide methanesulfonate, N-hydroxyglutarimide methane- 30 sulfonate, N-hydroxyglutarimide benzenesulfonate, N-hydroxyphthalimide methanesulfonate, N-hydroxyphthalimide benzenesulfonate, N-hydroxyphthalimide trifluoromethanesulfonate, N-hydroxyphthalimide p-toluenesulfonate, N-hydroxynaphthalimide methanesulfonate, N-hydroxynaphthalimide benzene- 35 sulfonate, N-hydroxy-5-norbornene-2,3-dicarboxyimide methanesulfonate, N-hydroxy-5-norbornene-2,3-dicarboxyimide

trifluoromethanesulfonate, and N-hydroxy-5-norbornene-2,3-dicarboxyimide p-toluenesulfonate.

Preferred among these photoacid generators are onium salts such as triphenylsulfonium trifluoromethanesulfonate, (p-tert-butoxyphenyl)diphenylsulfonium trifluoromethanesulfonate, tris(p-tert-butoxyphenyl)sulfonium trifluoromethanesulfonate, triphenylsulfonium p-toluenesulfonate, (p-tert-butoxyphenyl)diphenylsulfonium p-toluenesulfonate, tris(p-tert-butoxyphenyl)sulfonium p-toluenesulfonate, trinaphthylsulfonium trifluoromethanesulfonate, cyclohexylmethyl(2-oxocyclohexyl)sulfonium trifluoromethanesulfonate, (2-norbornyl)methyl(2-oxocyclohexyl)sulfonium trifluoromethanesulfonate, and 1,2'-naphthylcarbonylmethyltetrahydrothiophenium triflate; diazomethane derivatives such as bis(benzenesulfonyl)diazomethane, bis(p-toluenesulfonyl)diazomethane, bis(cyclohexylsulfonyl)diazomethane, bis(n-butylsulfonyl)diazomethane, bis(isobutylsulfonyl)diazomethane, bis(sec-butylsulfonyl)diazomethane, bis(n-propylsulfonyl)diazomethane, bis(isopropylsulfonyl)diazomethane, and bis(tert-butylsulfonyl)diazomethane; glyoxime derivatives such as bis-O-(p-toluenesulfonyl)- α -dimethylglyoxime and bis-O-(n-butan sulfonyl)- α -dimethylglyoxime; bissulfone derivatives such as bisnaphthylsulfonylmethane; and sulfonic acid esters of N-hydroxyimide compounds such as N-hydroxysuccinimide methanesulfonate, N-hydroxysuccinimide trifluoromethanesulfonate, N-hydroxysuccinimide 1-propanesulfonate, N-hydroxysuccinimide 2-propanesulfonate, N-hydroxysuccinimide 1-pentanesulfonate, N-hydroxysuccinimide p-toluenesulfonate, N-hydroxynaphthalimide methanesulfonate, and N-hydroxynaphthalimide benzenesulfonate.

These photoacid generators may be used singly or in combinations of two or more thereof. Onium salts are effective for improving rectangularity, while diazomethane derivatives and glyoxime derivatives are effective for reducing standing waves. The combination of an onium salt with a diazomethane or a glyoxime derivative allows for fine adjustment of the profile.

10073333 "024302
The photoacid generator is added in an amount of 0.1 to 15 parts, and especially 0.5 to 8 parts by weight, per 100 parts by weight of the base resin (all parts are by weight, hereinafter). Less than 0.1 part of the photoacid generator would provide a poor sensitivity whereas more than 15 parts of the photoacid generator would adversely affect transparency and resolution.

Organic solvent


The organic solvent used herein may be any organic solvent in which the base resin, photoacid generator, and other components are soluble. Illustrative, non-limiting, examples of the organic solvent include ketones such as cyclohexanone and methyl-2-n-amylketone; alcohols such as 3-methoxybutanol, 3-methyl-3-methoxybutanol, 1-methoxy-2-propanol, and 1-ethoxy-2-propanol; ethers such as propylene glycol monomethyl ether, ethylene glycol monomethyl ether, propylene glycol monoethyl ether, ethylene glycol monoethyl ether, propylene glycol dimethyl ether, and diethylene glycol dimethyl ether; esters such as propylene glycol monomethyl ether acetate, propylene glycol monoethyl ether acetate, ethyl lactate, ethyl pyruvate, butyl acetate, methyl 3-methoxypropionate, ethyl 3-ethoxypropionate, tert-butyl acetate, tert-butyl propionate, and propylene glycol mono-tert-butyl ether acetate; and lactones such as γ -butyrolactone. These solvents may be used alone or in combinations of two or more thereof. Of the above organic solvents, it is recommended to use diethylene glycol dimethyl ether and 1-ethoxy-2-propanol because the photoacid generator is most soluble therein, propylene glycol monomethyl ether acetate because it is a safe solvent, or a mixture thereof.

An appropriate amount of the organic solvent used is about 200 to 1,000 parts, especially about 400 to 800 parts by weight per 100 parts by weight of the base resin.

Other polymer

To the resist composition of the invention, another polymer other than the inventive polymer comprising

recurring units of formula (1) may also be added. The other polymers that can be added to the resist composition are, for example, those polymers comprising units of the following formula (R1) and/or (R2) and having a weight
 5 average molecular weight of about 1,000 to about 500,000, especially about 5,000 to about 100,000 although the other polymers are not limited thereto.



Herein, R^{001} is hydrogen, methyl or $CH_2CO_2R^{003}$. R^{002} is
 10 hydrogen, methyl or CO_2R^{003} . R^{003} is a straight, branched or cyclic alkyl group of 1 to 15 carbon atoms. R^{004} is hydrogen or a monovalent hydrocarbon group of 1 to 15 carbon atoms having a carboxyl or hydroxyl group. At least one of R^{005} to
 15 R^{008} represents a monovalent hydrocarbon group of 1 to 15 carbon atoms having a carboxyl or hydroxyl group while the

remaining R's independently represent hydrogen or a straight, branched or cyclic alkyl group of 1 to 15 carbon atoms.

Alternatively, R^{005} to R^{008} , taken together, may form a ring, and in that event, at least one of R^{005} to R^{008} is a divalent

hydrocarbon group of 1 to 15 carbon atoms having a carboxyl or hydroxyl group, while the remaining R's are independently single bonds or straight, branched or cyclic alkylene groups of 1 to 15 carbon atoms. R^{009} is a monovalent hydrocarbon

group of 2 to 15 carbon atoms containing at least one partial structure selected from among ether, aldehyde, ketone, ester, carbonate, acid anhydride, amide and imide.

At least one of R^{010} to R^{013} is a monovalent hydrocarbon group of 2 to 15 carbon atoms containing at least one partial structure selected from among ether, aldehyde, ketone, ester, carbonate, acid anhydride, amide and imide, while the

remaining R's are independently hydrogen or straight, branched or cyclic alkyl groups of 1 to 15 carbon atoms. R^{010} to R^{013} , taken together, may form a ring, and in that event, at least one of R^{010} to R^{013} is a divalent hydrocarbon group

of 1 to 15 carbon atoms containing at least one partial structure selected from among ether, aldehyde, ketone, ester, carbonate, acid anhydride, amide and imide, while the remaining R's are independently single bonds or straight, branched or cyclic alkylene groups of 1 to 15 carbon atoms.

R^{014} is a polycyclic hydrocarbon group having 7 to 15 carbon atoms or an alkyl group containing a polycyclic hydrocarbon group. R^{015} is an acid labile group. R^{016} is hydrogen or methyl. R^{017} is a straight, branched or cyclic alkyl group of 1 to 8 carbon atoms. X is CH_2 or an oxygen atom. Letter k'

is equal to 0 or 1; a_1' , a_2' , a_3' , b_1' , b_2' , b_3' , c_1' , c_2' , c_3' , d_1' , d_2' , d_3' , and e' are numbers from 0 to less than 1, satisfying

$a_1' + a_2' + a_3' + b_1' + b_2' + b_3' + c_1' + c_2' + c_3' + d_1' + d_2' + d_3' + e' = 1$; f' , g' , h' , i' , and j' are numbers from 0 to less than 1,

satisfying $f' + g' + h' + i' + j' = 1$; x' , y' and z' each are an integer of 0 to 3, satisfying $1 \leq x' + y' + z' \leq 5$ and $1 \leq y' + z' \leq 3$.

Exemplary groups of these R's are as exemplified above.

The inventive polymer (comprising recurring units of formula (1)) and the other polymer are preferably blended in a weight ratio from 100:0 to 10:90, more preferably from 100:0 to 20:80. If the blend ratio of the inventive polymer is below this range, the resist composition would become poor in some of the desired properties. The properties of the resist composition can be adjusted by properly changing the blend ratio of the inventive polymer.

The other polymer is not limited to one type and a mixture of two or more other polymers may be added. The use of plural polymers allows for easy adjustment of resist properties.


Dissolution regulator

To the resist composition, a dissolution regulator may be added. The dissolution regulator is a compound having on the molecule at least two phenolic hydroxyl groups, in which an average of from 0 to 100 mol% of all the hydrogen atoms on the phenolic hydroxyl groups are replaced with acid labile groups or a compound having on the molecule at least one carboxyl group, in which an average of 50 to 100 mol% of all the hydrogen atoms on the carboxyl groups are replaced with acid labile groups, both the compounds having an average molecular weight within a range of 100 to 1,000, and preferably 150 to 800.


The degree of substitution of the hydrogen atoms on the phenolic hydroxyl groups with acid labile groups is on average at least 0 mol%, and preferably at least 30 mol%, of all the phenolic hydroxyl groups. The upper limit is 100 mol%, and preferably 80 mol%. The degree of substitution of the hydrogen atoms on the carboxyl groups with acid labile groups is on average at least 50 mol%, and preferably at least 70 mol%, of all the carboxyl groups, with the upper limit being 100 mol%.

Preferable examples of such compounds having two or more phenolic hydroxyl groups or compounds having at least


one carboxyl group include those of formulas (D1) to (D14) below.




D1




D2




D3




D4




D5




D6




D7




D8




D9




D10




D11



D12



D13




D14


In these formulas, R^{201} and R^{202} are each hydrogen or a straight or branched alkyl or alkenyl of 1 to 8 carbon atoms; R^{203} is hydrogen, a straight or branched alkyl or alkenyl of 1 to 8 carbon atoms, or $-(R^{207})_h-COOH$; R^{204} is $-(CH_2)_i-$ (where $i = 2$ to 10), an arylene of 6 to 10 carbon atoms, carbonyl, sulfonyl, an oxygen atom, or a sulfur atom; R^{205} is an alkylene of 1 to 10 carbon atoms, an arylene of 6 to 10 carbon atoms, carbonyl, sulfonyl, an oxygen atom, or a sulfur atom; R^{206} is hydrogen, a straight or branched alkyl or alkenyl of 1 to 8 carbon atoms, or a hydroxyl-substituted phenyl or naphthyl; R^{207} is a straight or branched alkylene of 1 to 10 carbon atoms; R^{208} is hydrogen or hydroxyl; the letter j is an integer from 0 to 5; u and h are each 0 or 1; s , t , s' , t' , s'' , and t'' are each numbers which satisfy $s+t = 8$, $s'+t' = 5$, and $s''+t'' = 4$, and are such that each phenyl skeleton has at least one hydroxyl group; and α is a number such that the compounds of formula (D8) or (D9) have a molecular weight of from 100 to 1,000.

In the above formulas, suitable examples of R^{201} and R^{202} include hydrogen, methyl, ethyl, butyl, propyl, ethynyl, and cyclohexyl; suitable examples of R^{203} include the same groups as for R^{201} and R^{202} , as well as $-COOH$ and $-CH_2COOH$; suitable examples of R^{204} include ethylene, phenylene, carbonyl, sulfonyl, oxygen, and sulfur; suitable examples of R^{205} include methylene as well as the same groups as for R^{204} ; and suitable examples of R^{206} include hydrogen, methyl, ethyl, butyl, propyl, ethynyl, cyclohexyl, and hydroxyl-substituted phenyl or naphthyl.


Exemplary acid labile groups on the dissolution regulator include groups of the following general formulae (L1) to (L4), tertiary alkyl groups of 4 to 20 carbon atoms, trialkylsilyl groups in which each of the alkyls has 1 to 6 carbon atoms, and oxoalkyl groups of 4 to 20 carbon atoms.




(L1)



(L2)



(L3)



(L4)

In these formulas, R^{L01} and R^{L02} are each hydrogen or a straight, branched or cyclic alkyl having 1 to 18 carbon atoms; and R^{L03} is a monovalent hydrocarbon group of 1 to 18 carbon atoms which may contain a heteroatom (e.g., oxygen). A pair of R^{L01} and R^{L02} , a pair of R^{L01} and R^{L03} , or a pair of R^{L02} and R^{L03} may together form a ring, with the proviso that R^{L01} , R^{L02} , and R^{L03} are each a straight or branched alkylene of 1 to 18 carbon atoms when they form a ring. R^{L04} is a tertiary alkyl group of 4 to 20 carbon atoms, a trialksilyl group in which each of the alkyls has 1 to 6 carbon atoms, an oxoalkyl group of 4 to 20 carbon atoms, or a group of the formula (L1). R^{L05} is a monovalent hydrocarbon groups of 1 to 8 carbon atoms which may contain a hetero atom or a substituted or unsubstituted aryl group of 6 to 20 carbon atoms. R^{L06} is a monovalent hydrocarbon group of 1 to 8 carbon atoms which may contain a hetero atom or a substituted or unsubstituted aryl group of 6 to 20 carbon atoms. R^{L07} to R^{L16} independently represent hydrogen or monovalent hydrocarbon groups of 1 to 15 carbon atoms which may contain a hetero atom. Alternatively, R^{L07} to R^{L16} , taken together, may form a ring. Each of R^{L07} to R^{L16} represents a divalent $\text{C}_1\text{-C}_{15}$ hydrocarbon group which may contain a hetero atom, when they form a ring. Two of R^{L07} to R^{L16} which are attached to adjoining carbon atoms may bond together directly to form a double bond. Letter y is an integer of 0 to 6. Letter m is equal to 0 or 1, n is equal to 0, 1, 2 or 3, and $2m+n$ is equal to 2 or 3. Illustrative examples of these groups are as previously exemplified.

10073223-001300
The dissolution regulator may be formulated in an amount of 0 to 50 parts, preferably 0 to 40 parts, and more preferably 0 to 30 parts, per 100 parts of the base resin, and may be used singly or as a mixture of two or more
5 thereof. The use of more than 50 parts would lead to slimming of the patterned film, and thus a decline in resolution.

The dissolution regulator can be synthesized by introducing acid labile groups into a compound having
10 phenolic hydroxyl or carboxyl groups in accordance with an organic chemical formulation.

Basic compound

In the resist composition of the invention, a basic compound may be blended. A suitable basic compound used
15 herein is a compound capable of suppressing the rate of diffusion when the acid generated by the photoacid generator diffuses within the resist film. The inclusion of this type of basic compound holds down the rate of acid diffusion within the resist film, resulting in better resolution. In
20 addition, it suppresses changes in sensitivity following exposure, thus reducing substrate and environment dependence, as well as improving the exposure latitude and the pattern profile.

Examples of basic compounds include primary, secondary,
25 and tertiary aliphatic amines, mixed amines, aromatic amines, heterocyclic amines, carboxyl group-bearing nitrogenous compounds, sulfonyl group-bearing nitrogenous compounds, hydroxyl group-bearing nitrogenous compounds, hydroxyphenyl group-bearing nitrogenous compounds, alcoholic nitrogenous
30 compounds, amide derivatives, and imide derivatives.

Examples of suitable primary aliphatic amines include ammonia, methylamine, ethylamine, n-propylamine, isopropylamine, n-butylamine, iso-butylamine, sec-butylamine, tert-butylamine, pentylamine, tert-amylamine, cyclopentylamine,
35 hexylamine, cyclohexylamine, heptylamine, octylamine, nonylamine, decylamine, dodecylamine, cetylamine, methylenediamine, ethylenediamine, and tetraethylenepentamine.

Examples of suitable secondary aliphatic amines include dimethylamine, diethylamine, di-n-propylamine, di-iso-propylamine, di-n-butylamine, di-iso-butylamine, di-sec-butylamine, dipentylamine, dicyclopentylamine, dihexylamine, dicyclohexylamine, diheptylamine, dioctylamine, dinonylamine, didecylamine, didodecylamine, dicetylamine, N,N-dimethylmethylenediamine, N,N-dimethylethylenediamine, and N,N-dimethyltetraethylenepentamine. Examples of suitable tertiary aliphatic amines include trimethylamine, triethylamine, tri-n-propylamine, tri-iso-propylamine, tri-n-butylamine, tri-iso-butylamine, tri-sec-butylamine, tripentylamine, tricyclopentylamine, trihexylamine, tricyclohexylamine, triheptylamine, trioctylamine, trinonylamine, tridecylamine, tridodecylamine, tricetylamine, N,N,N',N'-tetramethylmethylenediamine, N,N,N',N'-tetramethylethylenediamine, and N,N,N',N'-tetramethyltetraethylenepentamine.

Examples of suitable mixed amines include dimethylethylamine, methylethylpropylamine, benzylamine, phenethylamine, and benzyldimethylamine. Examples of suitable aromatic and heterocyclic amines include aniline derivatives (e.g., aniline, N-methylaniline, N-ethylaniline, N-propylaniline, N,N-dimethylaniline, 2-methylaniline, 3-methylaniline, 4-methylaniline, ethylaniline, propylaniline, trimethylaniline, 2-nitroaniline, 3-nitroaniline, 4-nitroaniline, 2,4-dinitroaniline, 2,6-dinitroaniline, 3,5-dinitroaniline, and N,N-dimethyltoluidine), diphenyl(p-tolyl)amine, methyldiphenylamine, triphenylamine, phenylenediamine, naphthylamine, diaminonaphthalene, pyrrole derivatives (e.g., pyrrole, 2H-pyrrole, 1-methylpyrrole, 2,4-dimethylpyrrole, 2,5-dimethylpyrrole, and N-methylpyrrole), oxazole derivatives (e.g., oxazole and isooxazole), thiazole derivatives (e.g., thiazole and isothiazole), imidazole derivatives (e.g., imidazole, 4-methylimidazole, and 4-methyl-2-phenylimidazole), pyrazole derivatives, furazan derivatives, pyrroline derivatives (e.g., pyrroline and 2-methyl-1-pyrroline), pyrrolidine derivatives (e.g., pyrrolidine, N-methylpyrrolidine, pyrrolidinone, and N-

methylpyrrolidone), imidazoline derivatives, imidazolidine
 derivatives, pyridine derivatives (e.g., pyridine, methyl-
 pyridine, ethylpyridine, propylpyridine, butylpyridine, 4-
 (1-butylpentyl)pyridine, dimethylpyridine, trimethylpyridine,
 5 triethylpyridine, phenylpyridine, 3-methyl-2-phenylpyridine,
 4-tert-butylpyridine, diphenylpyridine, benzylpyridine,
 methoxypyridine, butoxypyridine, dimethoxypyridine, 1-
 methyl-2-pyridone, 4-pyrrolidinopyridine, 1-methyl-4-
 phenylpyridine, 2-(1-ethylpropyl)pyridine, aminopyridine,
 10 and dimethylaminopyridine), pyridazine derivatives,
 pyrimidine derivatives, pyrazine derivatives, pyrazoline
 derivatives, pyrazolidine derivatives, piperidine
 derivatives, piperazine derivatives, morpholine derivatives,
 indole derivatives, isoindole derivatives, 1H-indazole
 15 derivatives, indoline derivatives, quinoline derivatives
 (e.g., quinoline and 3-quinolinecarbonitrile), isoquinoline
 derivatives, cinnoline derivatives, quinazoline derivatives,
 quinoxaline derivatives, phthalazine derivatives, purine
 derivatives, pteridine derivatives, carbazole derivatives,
 20 phenanthridine derivatives, acridine derivatives, phenazine
 derivatives, 1,10-phenanthroline derivatives, adenine
 derivatives, adenosine derivatives, guanine derivatives,
 guanosine derivatives, uracil derivatives, and uridine
 derivatives.

25 Examples of suitable carboxyl group-bearing
 nitrogenous compounds include aminobenzoic acid,
 indolecarboxylic acid, and amino acid derivatives (e.g.
 nicotinic acid, alanine, alginine, aspartic acid, glutamic
 acid, glycine, histidine, isoleucine, glycyllleucine, leucine,
 30 methionine, phenylalanine, threonine, lysine, 3-
 aminopyrazine-2-carboxylic acid, and methoxyalanine).
 Examples of suitable sulfonyl group-bearing nitrogenous
 compounds include 3-pyridinesulfonic acid and pyridinium p-
 toluenesulfonate. Examples of suitable hydroxyl group-
 35 bearing nitrogenous compounds, hydroxyphenyl group-bearing
 nitrogenous compounds, and alcoholic nitrogenous compounds
 include 2-hydroxypyridine, aminocresol, 2,4-quinolinediol,


3-indolemethanol hydrate, monoethanolamine, diethanolamine, triethanolamine, N-ethyldiethanolamine, N,N-diethylethanolamine, triisopropanolamine, 2,2'-iminodiethanol, 2-aminoethanol, 3-amino-1-propanol, 4-amino-1-butanol, 4-(2-hydroxyethyl)morpholine, 2-(2-hydroxyethyl)pyridine, 1-(2-hydroxyethyl)piperazine, 1-[2-(2-hydroxyethoxy)ethyl]-piperazine, piperidine ethanol, 1-(2-hydroxyethyl)-pyrrolidine, 1-(2-hydroxyethyl)-2-pyrrolidinone, 3-piperidino-1,2-propanediol, 3-pyrrolidino-1,2-propanediol, 8-hydroxyjulolidine, 3-quinuclidinol, 3-tropanol, 1-methyl-2-pyrrolidine ethanol, 1-aziridine ethanol, N-(2-hydroxyethyl)phthalimide, and N-(2-hydroxyethyl)isonicotinamide.

Examples of suitable amide derivatives include formamide, N-methylformamide, N,N-dimethylformamide, acetamide, N-methylacetamide, N,N-dimethylacetamide, propionamide, and benzamide. Suitable imide derivatives include phthalimide, succinimide, and maleimide.


In addition, basic compounds of the following general formula (B1) may also be included alone or in admixture.




In the formula, n is equal to 1, 2 or 3; Y is independently hydrogen or a straight, branched or cyclic alkyl group of 1 to 20 carbon atoms which may contain a hydroxyl group or ether; and X is independently selected from groups of the following general formulas (X1) to (X3), and two or three X's may bond together to form a ring.



(X1)



(X2)



(X3)

In the formulas, R^{300} , R^{302} and R^{305} are independently straight or branched alkylene groups of 1 to 4 carbon atoms; R^{301} and R^{304} are independently hydrogen, straight, branched or cyclic alkyl groups of 1 to 20 carbon atoms, which may contain at least one hydroxyl group, ether, ester or lactone

ring; R³⁰³ is a single bond or a straight or branched
alkylene group of 1 to 4 carbon atoms; and R³⁰⁶ is a single
bond or a straight, branched or cyclic alkyl group of 1 to
20 carbon atoms which may contain one or more hydroxyl
groups, ether groups, ester groups or lactone rings.

Illustrative examples of the compounds of formula (B1)
include tris(2-methoxymethoxyethyl)amine, tris{2-(methoxy-
ethoxy)ethyl}amine, tris{2-(2-methoxyethoxymethoxy)ethyl}-
amine, tris{2-(1-methoxyethoxy)ethyl}amine, tris{2-(1-
ethoxyethoxy)ethyl}amine, tris{2-(1-ethoxypropoxy)ethyl}-
amine, tris[2-{2-(2-hydroxyethoxy)ethoxy}ethyl]amine,
4,7,13,16,21,24-hexaoxa-1,10-diazabicyclo[8.8.8]hexacosane,
4,7,13,18-tetraoxa-1,10-diazabicyclo[8.5.5]eicosane,
1,4,10,13-tetraoxa-7,16-diazabicyclooctadecane, 1-aza-12-
crown-4, 1-aza-15-crown-5, 1-aza-18-crown-6, tris(2-
formyloxyethyl)amine, tris(2-acetoxyethyl)amine, tris(2-
propionyloxyethyl)amine, tris(2-butyryloxyethyl)amine,
tris(2-isobutyryloxyethyl)amine, tris(2-valeryloxyethyl)-
amine, tris(2-pivaloyloxyethyl)amine, N,N-bis(2-acetoxy-
ethyl)-2-(acetoxyacetoxy)ethylamine, tris(2-methoxycarbonyl-
oxyethyl)amine, tris(2-tert-butoxycarbonyloxyethyl)amine,
tris[2-(2-oxopropoxy)ethyl]amine, tris[2-(methoxycarbonyl-
methyl)oxyethyl]amine, tris[2-(tert-butoxycarbonylmethyl-
oxy)ethyl]amine, tris[2-(cyclohexyloxycarbonylmethyloxy)-
ethyl]amine, tris(2-methoxycarbonylethyl)amine, tris(2-
ethoxycarbonylethyl)amine, N,N-bis(2-hydroxyethyl)-2-
(methoxycarbonyl)ethylamine, N,N-bis(2-acetoxyethyl)-2-
(methoxycarbonyl)ethylamine, N,N-bis(2-hydroxyethyl)-2-
(ethoxycarbonyl)ethylamine, N,N-bis(2-acetoxyethyl)-2-
(ethoxycarbonyl)ethylamine, N,N-bis(2-hydroxyethyl)-2-(2-
methoxyethoxycarbonyl)ethylamine, N,N-bis(2-acetoxyethyl)-2-
(2-methoxyethoxycarbonyl)ethylamine, N,N-bis(2-hydroxy-
ethyl)-2-(2-hydroxyethoxycarbonyl)ethylamine, N,N-bis(2-
acetoxyethyl)-2-(2-acetoxyethoxycarbonyl)ethylamine, N,N-
bis(2-hydroxyethyl)-2-[(methoxycarbonyl)methoxycarbonyl]-
ethylamine, N,N-bis(2-acetoxyethyl)-2-[(methoxycarbonyl)-
methoxycarbonyl]ethylamine, N,N-bis(2-hydroxyethyl)-2-(2-

oxopropoxycarbonyl)ethylamine, N,N-bis(2-acetoxyethyl)-2-(2-oxopropoxycarbonyl)ethylamine, N,N-bis(2-hydroxyethyl)-2-(tetrahydrofurfuryloxycarbonyl)ethylamine, N,N-bis(2-acetoxyethyl)-2-(tetrahydrofurfuryloxycarbonyl)ethylamine, N,N-bis(2-hydroxyethyl)-2-[(2-oxotetrahydrofuran-3-yl)oxycarbonyl]ethylamine, N,N-bis(2-acetoxyethyl)-2-[(2-oxotetrahydrofuran-3-yl)oxycarbonyl]ethylamine, N,N-bis(2-hydroxyethyl)-2-(4-hydroxybutoxycarbonyl)ethylamine, N,N-bis(2-formyloxyethyl)-2-(4-formyloxybutoxycarbonyl)ethylamine, N,N-bis(2-formyloxyethyl)-2-(2-formyloxyethoxycarbonyl)-ethylamine, N,N-bis(2-methoxyethyl)-2-(methoxycarbonyl)-ethylamine, N-(2-hydroxyethyl)-bis[2-(methoxycarbonyl)-ethyl]amine, N-(2-acetoxyethyl)-bis[2-(methoxycarbonyl)-ethyl]amine, N-(2-hydroxyethyl)-bis[2-(ethoxycarbonyl)-ethyl]amine, N-(2-acetoxyethyl)-bis[2-(ethoxycarbonyl)-ethyl]amine, N-(3-hydroxy-1-propyl)-bis[2-(methoxycarbonyl)-ethyl]amine, N-(3-acetoxy-1-propyl)-bis[2-(methoxycarbonyl)-ethyl]amine, N-(2-methoxyethyl)-bis[2-(methoxycarbonyl)-ethyl]amine, N-butyl-bis[2-(methoxycarbonyl)ethyl]amine, N-butyl-bis[2-(2-methoxyethoxycarbonyl)ethyl]amine, N-methyl-bis(2-acetoxyethyl)amine, N-ethyl-bis(2-acetoxyethyl)amine, N-methyl-bis(2-pivaloyloxyethyl)amine, N-ethyl-bis[2-(methoxycarbonyloxy)ethyl]amine, N-ethyl-bis[2-(tert-butoxycarbonyloxy)ethyl]amine, tris(methoxycarbonylmethyl)amine, tris(ethoxycarbonylmethyl)amine, N-butyl-bis(methoxycarbonylmethyl)amine, N-hexyl-bis(methoxycarbonylmethyl)amine, and β -(diethylamino)- δ -valerolactone.

Also useful are one or more of cyclic structure-bearing basic compounds having the following general formula (B2).




Herein X is as defined above, and R^{307} is a straight or branched alkylene group of 2 to 20 carbon atoms which may

contain one or more carbonyl groups, ether structures, ester structures or sulfide structures.


Illustrative examples of the cyclic structure-bearing basic compounds having formula (B2) include

5 1-[2-(methoxymethoxy)ethyl]pyrrolidine, 1-[2-(methoxy-
methoxy)ethyl]piperidine, 4-[2-(methoxymethoxy)ethyl]-
morpholine, 1-[2-[(2-methoxyethoxy)methoxy]ethyl]pyrrolidine,
1-[2-[(2-methoxyethoxy)methoxy]ethyl]piperidine, 4-[2-[(2-
methoxyethoxy)methoxy]ethyl]morpholine, 2-(1-pyrrolidinyl)-
10 ethyl acetate, 2-piperidinoethyl acetate, 2-morpholinoethyl
acetate, 2-(1-pyrrolidinyl)ethyl formate, 2-piperidinoethyl
propionate, 2-morpholinoethyl acetoxyacetate, 2-(1-
pyrrolidinyl)ethyl methoxyacetate, 4-[2-(methoxycarbonyl-
oxy)ethyl]morpholine, 1-[2-(t-butoxycarbonyloxy)ethyl]-
15 piperidine, 4-[2-(2-methoxyethoxycarbonyloxy)ethyl]-
morpholine, methyl 3-(1-pyrrolidinyl)propionate, methyl 3-
piperidinopropionate, methyl 3-morpholinopropionate, methyl
3-(thiomorpholino)propionate, methyl 2-methyl-3-(1-
pyrrolidinyl)propionate, ethyl 3-morpholinopropionate,
20 methoxycarbonylmethyl 3-piperidinopropionate, 2-hydroxyethyl
3-(1-pyrrolidinyl)propionate, 2-acetoxyethyl 3-morpholino-
propionate, 2-oxotetrahydrofuran-3-yl 3-(1-pyrrolidinyl)-
propionate, tetrahydrofurfuryl 3-morpholinopropionate,
glycidyl 3-piperidinopropionate, 2-methoxyethyl 3-
25 morpholinopropionate, 2-(2-methoxyethoxy)ethyl 3-(1-
pyrrolidinyl)propionate, butyl 3-morpholinopropionate,
cyclohexyl 3-piperidinopropionate, α -(1-pyrrolidinyl)methyl-
 γ -butyrolactone, β -piperidino- γ -butyrolactone, β -morpholino-
 δ -valerolactone, methyl 1-pyrrolidinylacetate, methyl
30 piperidinoacetate, methyl morpholinoacetate, methyl
thiomorpholinoacetate, ethyl 1-pyrrolidinylacetate, and 2-
methoxyethyl morpholinoacetate.


Also, one or more of cyano-bearing basic compounds
having the following general formulae (B3) to (B6) may be
35 blended.




(B3)



(B4)



(B5)



(B6)

Herein, X, R^{307} and n are as defined above, and R^{308} and R^{309} each are independently a straight or branched alkylene group of 1 to 4 carbon atoms.

5

Illustrative examples of the cyano-bearing basic

compounds having formulae (B3) to (B6) include

3-(diethylamino)propiononitrile, N,N-bis(2-hydroxyethyl)-3-aminopropiononitrile, N,N-bis(2-acetoxyethyl)-3-aminopropiononitrile, N,N-bis(2-formyloxyethyl)-3-aminopropiononitrile, N,N-bis(2-methoxyethyl)-3-aminopropiononitrile, N,N-bis[2-(methoxymethoxy)ethyl]-3-aminopropiononitrile, methyl N-(2-cyanoethyl)-N-(2-methoxyethyl)-3-aminopropionate, methyl N-(2-cyanoethyl)-N-(2-hydroxyethyl)-3-aminopropionate, methyl N-(2-acetoxyethyl)-N-(2-cyanoethyl)-3-aminopropionate, N-(2-cyanoethyl)-N-ethyl-3-aminopropiononitrile, N-(2-cyanoethyl)-N-(2-hydroxyethyl)-3-aminopropiononitrile, N-(2-acetoxyethyl)-N-(2-cyanoethyl)-3-aminopropiononitrile, N-(2-cyanoethyl)-N-(2-formyloxyethyl)-3-aminopropiononitrile, N-(2-cyanoethyl)-N-(2-methoxyethyl)-3-aminopropiononitrile, N-(2-cyanoethyl)-N-[2-(methoxymethoxy)ethyl]-3-aminopropiononitrile, N-(2-cyanoethyl)-N-(3-hydroxy-1-propyl)-3-aminopropiononitrile, N-(3-acetoxy-1-propyl)-N-(2-cyanoethyl)-3-aminopropiononitrile, N-(2-cyanoethyl)-N-(3-formyloxy-1-propyl)-3-aminopropiononitrile, N-(2-cyanoethyl)-N-tetrahydrofurfuryl-3-aminopropiononitrile, N,N-bis(2-cyanoethyl)-3-aminopropiononitrile, diethylaminoacetone nitrile, N,N-bis(2-hydroxyethyl)aminoacetone nitrile, N,N-bis(2-acetoxyethyl)aminoacetone nitrile, N,N-bis(2-formyloxyethyl)aminoacetone nitrile, N,N-bis(2-methoxyethyl)aminoacetone nitrile, N,N-bis[2-(methoxymethoxy)ethyl]aminoacetone nitrile, methyl N-cyanomethyl-N-(2-methoxyethyl)-3-aminopropionate, methyl N-cyanomethyl-N-(2-hydroxyethyl)-3-aminopropionate, methyl N-

(2-acetoxyethyl)-N-cyanomethyl-3-aminopropionate, N-cyanomethyl-N-(2-hydroxyethyl)aminoacetonitrile, N-(2-acetoxyethyl)-N-(cyanomethyl)aminoacetonitrile, N-cyanomethyl-N-(2-formyloxyethyl)aminoacetonitrile, N-cyanomethyl-N-(2-methoxyethyl)aminoacetonitrile, N-cyanomethyl-N-[2-(methoxymethoxy)ethyl]aminoacetonitrile, N-cyanomethyl-N-(3-hydroxy-1-propyl)aminoacetonitrile, N-(3-acetoxy-1-propyl)-N-(cyanomethyl)aminoacetonitrile, N-cyanomethyl-N-(3-formyloxy-1-propyl)aminoacetonitrile, N,N-bis(cyanomethyl)aminoacetonitrile, 1-pyrrolidinepropiononitrile, 1-piperidinepropiononitrile, 4-morpholinepropiononitrile, 1-pyrrolidineacetonitrile, 1-piperidineacetonitrile, 4-morpholineacetonitrile, cyanomethyl 3-diethylaminopropionate, cyanomethyl N,N-bis(2-hydroxyethyl)-3-aminopropionate, cyanomethyl N,N-bis(2-acetoxyethyl)-3-aminopropionate, cyanomethyl N,N-bis(2-formyloxyethyl)-3-aminopropionate, cyanomethyl N,N-bis(2-methoxyethyl)-3-aminopropionate, cyanomethyl N,N-bis[2-(methoxymethoxy)ethyl]-3-aminopropionate, 2-cyanoethyl 3-diethylaminopropionate, 2-cyanoethyl N,N-bis(2-hydroxyethyl)-3-aminopropionate, 2-cyanoethyl N,N-bis(2-acetoxyethyl)-3-aminopropionate, 2-cyanoethyl N,N-bis(2-formyloxyethyl)-3-aminopropionate, 2-cyanoethyl N,N-bis(2-methoxyethyl)-3-aminopropionate, 2-cyanoethyl N,N-bis[2-(methoxymethoxy)ethyl]-3-aminopropionate, cyanomethyl 1-pyrrolidinepropionate, cyanomethyl 1-piperidinepropionate, cyanomethyl 4-morpholinepropionate, 2-cyanoethyl 1-pyrrolidinepropionate, 2-cyanoethyl 1-piperidinepropionate, and 2-cyanoethyl 4-morpholinepropionate.

The basic compound is preferably formulated in an amount of 0.001 to 10 parts, and especially 0.01 to 1 part, per part of the photoacid generator. Less than 0.001 part of the basic compound may fail to achieve the desired effects thereof, while the use of more than 10 parts would result in too low a sensitivity and resolution.

Other components

In the resist composition, a compound bearing a $\equiv\text{C}-\text{COOH}$ group in a molecule may be blended. Exemplary, non-


limiting compounds bearing a $\equiv\text{C}-\text{COOH}$ group include one or more compounds selected from Groups I and II below.

Including this compound improves the PED stability of the resist and ameliorates edge roughness on nitride film


5 substrates.

Group I:


Compounds in which some or all of the hydrogen atoms on the phenolic hydroxyl groups of the compounds of general formulas (A1) to (A10) below have been replaced with $-\text{R}^{401}-$
 10 COOH (wherein R^{401} is a straight or branched alkylene of 1 to 10 carbon atoms), and in which the molar ratio $\text{C}/(\text{C}+\text{D})$ of phenolic hydroxyl groups (C) to $\equiv\text{C}-\text{COOH}$ groups (D) in the molecule is from 0.1 to 1.0.




(A1)




(A2)




(A3)




(A4)




(A5)




(A6)




(A7)



(A8)



(A9)




(A10)

15 In these formulas, R^{408} is hydrogen or methyl; R^{402} and R^{403} are each hydrogen or a straight or branched alkyl or


alkenyl of 1 to 8 carbon atoms; R^{404} is hydrogen, a straight or branched alkyl or alkenyl of 1 to 8 carbon atoms, or a $-(R^{409})_h-COOR'$ group (R' being hydrogen or $-R^{409}-COOH$); R^{405} is $-(CH_2)_i-$ (wherein i is 2 to 10), an arylene of 6 to 10 carbon atoms, carbonyl, sulfonyl, an oxygen atom, or a sulfur atom; R^{406} is an alkylene of 1 to 10 carbon atoms, an arylene of 6 to 10 carbon atoms, carbonyl, sulfonyl, an oxygen atom, or a sulfur atom; R^{407} is hydrogen, a straight or branched alkyl or alkenyl of 1 to 8 carbon atoms, or a hydroxyl-substituted phenyl or naphthyl; R^{409} is a straight or branched alkylene of 1 to 10 carbon atoms; R^{410} is hydrogen, a straight or branched alkyl or alkenyl of 1 to 8 carbon atoms, or a $-R^{411}-COOH$ group; R^{411} is a straight or branched alkylene of 1 to 10 carbon atoms; the letter j is an integer from 0 to 5; u and h are each 0 or 1; $s_1, t_1, s_2, t_2, s_3, t_3, s_4,$ and t_4 are each numbers which satisfy $s_1+t_1 = 8, s_2+t_2 = 5, s_3+t_3 = 4,$ and $s_4+t_4 = 6,$ and are such that each phenyl skeleton has at least one hydroxyl group; κ is a number such that the compound of formula (A6) may have a weight average molecular weight of 1,000 to 5,000; and λ is a number such that the compound of formula (A7) may have a weight average molecular weight of 1,000 to 10,000.

Group II:


Compounds of general formulas (A11) to (A15) below.




(A11)




(A12)



(A13)




(A14)




(A15)

In these formulas, R^{402} , R^{403} , and R^{411} are as defined above; R^{412} is hydrogen or hydroxyl; s_5 and t_5 are numbers which satisfy $s_5 \geq 0$, $t_5 \geq 0$, and $s_5 + t_5 = 5$; and h' is equal to 0 or 1.


- 5 Illustrative, non-limiting examples of the compound bearing a $\equiv C-COOH$ group include compounds of the general formulas AI-1 to AI-14 and AII-1 to AII-10 below.




AI-1




AI-2




AI-3




AI-4




AI-5




AI-6




AI-7




AI-8




AI-9




AI-10




AI-11



AI-12




AI-13




AI-14


In the above formulas, R'' is hydrogen or a CH₂COOH group such that the CH₂COOH group accounts for 10 to 100 mol% of R'' in each compound, α and κ are as defined above.




AII-1




AII-2




AII-3




AII-4




AII-5




AII-6




AII-7



AII-8



AII-9




AII-10


The compound bearing a $\equiv\text{C}-\text{COOH}$ group within the molecule may be used singly or as combinations of two or more thereof.

The compound bearing a $\equiv\text{C}-\text{COOH}$ group within the molecule is added in an amount ranging from 0 to 5 parts, preferably 0.1 to 5 parts, more preferably 0.1 to 3 parts, further preferably 0.1 to 2 parts, per 100 parts of the base resin. More than 5 parts of the compound can reduce the resolution of the resist composition.

The resist composition of the invention may additionally include an acetylene alcohol derivative for the purpose of enhancing the shelf stability. Preferred acetylene alcohol derivatives are those having the general formula (S1) or (S2) below.



(S1)



(S2)

In the formulas, R^{501} , R^{502} , R^{503} , R^{504} , and R^{505} are each hydrogen or a straight, branched, or cyclic alkyl of 1 to 8 carbon atoms; and X and Y are each 0 or a positive number, satisfying $0 \leq X \leq 30$, $0 \leq Y \leq 30$, and $0 \leq X+Y \leq 40$.

Preferable examples of the acetylene alcohol derivative include Surfynol 61, Surfynol 82, Surfynol 104, Surfynol 104E, Surfynol 104H, Surfynol 104A, Surfynol TG, Surfynol PC, Surfynol 440, Surfynol 465, and Surfynol 485 from Air Products and Chemicals Inc., and Surfynol E1004 from Nisshin Chemical Industry K.K.

The acetylene alcohol derivative is preferably added in an amount of 0.01 to 2% by weight, and more preferably 0.02 to 1% by weight, per 100% by weight of the resist composition. Less than 0.01% by weight would be ineffective for improving coating characteristics and shelf stability, whereas more than 2% by weight would result in a resist having a low resolution.

The resist composition of the invention may include optional ingredients, for example, a surfactant which is commonly used for improving the coating characteristics. Optional ingredients may be added in conventional amounts so long as this does not compromise the objects of the invention.

Nonionic surfactants are preferred, examples of which include perfluoroalkylpolyoxyethylene ethanols, fluorinated alkyl esters, perfluoroalkylamine oxides, perfluoroalkyl EO-addition products, and fluorinated organosiloxane compounds. Useful surfactants are commercially available under the trade names Florade FC-430 and FC-431 from Sumitomo 3M, Ltd., Surflon S-141 and S-145, KH-10, KH-20, KH-30 and KH-40 from Asahi Glass Co., Ltd., Unidyne DS-401, DS-403 and DS-451 from Daikin Industry Co., Ltd., Megaface F-8151 from Dai-Nippon Ink & Chemicals, Inc., and X-70-092 and X-70-093 from Shin-Etsu Chemical Co., Ltd. Preferred surfactants are Florade FC-430 from Sumitomo 3M, Ltd., KH-20 and KH-30 from Asahi Glass Co., Ltd., and X-70-093 from Shin-Etsu Chemical Co., Ltd.

Pattern formation using the resist composition of the invention may be carried out by a known lithographic technique. For example, the resist composition is applied onto a substrate such as a silicon wafer by spin coating or the like to form a resist film having a thickness of 0.2 to 2.0 μm , which is then pre-baked on a hot plate at 60 to 150°C for 1 to 10 minutes, and preferably at 80 to 130°C for 1 to 5 minutes. A patterning mask having the desired pattern is then placed over the resist film, and the film exposed through the mask to an electron beam or to high-energy radiation such as deep-UV rays, an excimer laser, or x-rays in a dose of about 1 to 200 mJ/cm^2 , and preferably about 5 to 100 mJ/cm^2 , then post-exposure baked (PEB) on a hot plate at 60 to 150°C for 1 to 5 minutes, and preferably at 80 to 130°C for 1 to 3 minutes. Finally, development is carried out using as the developer an aqueous alkali solution, such as a 0.1 to 5% (preferably 2 to 3%) aqueous

10073223-027302
solution of tetramethylammonium hydroxide (TMAH), this being
done by a conventional method such as dipping, puddling, or
spraying for a period of 0.1 to 3 minutes, and preferably
0.5 to 2 minutes. These steps result in the formation of
5 the desired pattern on the substrate. Of the various types
of high-energy radiation that may be used, the resist
composition of the invention is best suited to fine pattern
formation with, in particular, deep-UV rays having a
wavelength of 248 to 193 nm, an excimer laser, x-rays, or an
10 electron beam. The desired pattern may not be obtainable
outside the upper and lower limits of the above range.

The resist composition comprising the inventive
polymer as a base resin lends itself to micropatterning with
electron beams or deep-UV rays since it is sensitive to
15 high-energy radiation and has excellent sensitivity,
resolution, and etching resistance. Especially because of
the minimized absorption at the exposure wavelength of an
ArF or KrF excimer laser, a finely defined pattern having
sidewalls perpendicular to the substrate can easily be
20 formed.

EXAMPLE

Synthesis Examples and Examples are given below by way
of illustration and not by way of limitation. The
25 abbreviation Mw is a weight average molecular weight as
measured by GPC using a polystyrene standard, and SEM is
scanning electron microscope.

Polymers within the scope of the invention were
synthesized by the following procedure.

Synthesis Example 1

Synthesis of Polymer 1

A mixture of 16.00 g (160 mmol) of isobutyl vinyl
ether, 14.88 g (60 mmol) of 2-ethyl-2-adamantyl
methacrylate, and 3.43 g of 1,4-dioxane was heated at 60°C.
35 To the solution was added 0.74 g (3 mmol) of 2,2'-
azobis(2,4-dimethylvaleronitrile). The solution was stirred
for 15 hours while keeping at 60°C. The reaction solution

was cooled to room temperature and dissolved in 50 ml of acetone, which with vigorous stirring, was added dropwise to 1000 ml of isopropyl alcohol. The resulting solids were collected by filtration and dried in vacuum at 40°C for 15 hours, obtaining 11.28 g of a white powder. On ¹H-NMR analysis, it was confirmed to be a polymer designated Polymer 1 whose structure is shown below.

Synthesis Examples 2-12

Synthesis of Polymers 2-12

Polymers 2 to 12 were synthesized by the same procedure as above or a well-known procedure.

Synthesis Example 13

Synthesis of Polymer 13

A mixture of 22.40 g (160 mmol) of 5-methoxymethyl-7-oxabicyclo[2.2.1]hept-2-ene, 14.88 g (60 mmol) of 2-ethyl-2-adamantyl methacrylate, and 5.33 g of 1,4-dioxane was heated at 60°C. To the solution was added 0.74 g (3 mmol) of 2,2'-azobis(2,4-dimethylvaleronitrile). The solution was stirred for 15 hours while keeping at 60°C. The reaction solution was cooled to room temperature and dissolved in 50 ml of acetone, which with vigorous stirring, was added dropwise to 1000 ml of isopropyl alcohol. The resulting solids were collected by filtration and dried in vacuum at 40°C for 15 hours, obtaining 11.26 g of a white powder. On ¹H-NMR analysis, it was confirmed to be a polymer designated Polymer 13 whose structure is shown below.

Synthesis Examples 14-20

Synthesis of Polymers 14-20

Polymers 14 to 20 were synthesized by the same procedure as above or a well-known procedure.

Synthesis Example 21

Synthesis of Polymer 21


A mixture of 20.80 g (160 mmol) of acrolein diethyl acetal, 14.88 g (60 mmol) of 2-ethyl-2-adamantyl methacrylate, and 3.57 g of 1,4-dioxane was heated at 60°C. To the solution was added 0.74 g (3 mmol) of 2,2'-azobis(2,4-dimethylvaleronitrile). The solution was stirred for 15 hours while keeping at 60°C. The reaction solution was cooled to room temperature and dissolved in 50 ml of acetone, which with vigorous stirring, was added dropwise to 1000 ml of isopropyl alcohol. The resulting solids were collected by filtration and dried in vacuum at 40°C for 15 hours, obtaining 10.08 g of a white powder. On ¹H-NMR analysis, it was confirmed to be a polymer designated Polymer 21 whose structure is shown below.

Synthesis Examples 22-28


Synthesis of Polymers 22-28

Polymers 22 to 28 were synthesized by the same procedure as above or a well-known procedure.



(Polymer 1)
(x = 0.40, d = 0.60, Mw = 6,800)





(Polymer 2)
(x = 0.40, d = 0.60, Mw = 7,300)





(Polymer 3)
($x = 0.40$, $d = 0.60$, $M_w = 8,000$)





(Polymer 4)
($x = 0.40$, $d = 0.60$, $M_w = 6,100$)





(Polymer 5)
($x = 0.40$, $d = 0.60$, $M_w = 6,800$)




(Polymer 6)
($x = 0.40$, $d = 0.60$, $M_w = 7,300$)




(Polymer 7)
($x = 0.40$, $d = 0.60$, $M_w = 8,000$)




(Polymer 8)
($x = 0.40$, $d = 0.60$, $M_w = 6,100$)




(Polymer 9)
($x = 0.40$, $d = 0.60$, $M_w = 7,400$)




(Polymer 10)
($x = 0.40$, $d = 0.60$, $M_w = 7,100$)




(Polymer 11)
($x = 0.40$, $a = 0.10$, $d = 0.50$,
 $M_w = 8,600$)




(Polymer 12)
($x = 0.30$, $b = 0.20$, $d = 0.50$,
 $M_w = 8,200$)




(Polymer 13)
($x = 0.40$, $d = 0.60$, $M_w = 7,700$)




(Polymer 14)
($x = 0.40$, $d = 0.60$, $M_w = 8,100$)




(Polymer 15)
($x = 0.40$, $d = 0.60$, $M_w = 7,200$)




(Polymer 16)
($x = 0.40$, $d = 0.60$, $M_w = 7,500$)




(Polymer 17)
($x = 0.40$, $d = 0.60$, $M_w = 8,800$)




(Polymer 18)
($x = 0.40$, $d = 0.60$, $M_w = 8,500$)




(Polymer 19)
($x = 0.40$, $a = 0.10$, $d = 0.50$,
 $M_w = 8,600$)




(Polymer 20)
($x = 0.30$, $b = 0.20$, $d = 0.50$,
 $M_w = 8,300$)




(Polymer 21)
($x = 0.40$, $d = 0.60$, $M_w = 6,800$)




(Polymer 22)
($x = 0.40$, $d = 0.60$, $M_w = 7,300$)




(Polymer 23)
($x = 0.40$, $d = 0.60$, $M_w = 8,000$)




(Polymer 24)
($x = 0.40$, $d = 0.60$, $M_w = 6,100$)




(Polymer 25)
($x = 0.40$, $d = 0.60$, $M_w = 7,400$)




(Polymer 26)
($x = 0.40$, $d = 0.60$, $M_w = 7,100$)



(Polymer 27)
($x = 0.40$, $a = 0.10$, $d = 0.50$,
 $M_w = 8,600$)



(Polymer 28)
 (x = 0.30, b = 0.20, d = 0.50,
 Mw = 8,200)




Resist compositions were formulated using inventive polymers and examined for resolution upon ArF excimer laser exposure.


5 Examples 1 to 55

Evaluation of resist resolution


Resist compositions were prepared by using Polymers 1 to 28 as the base resin, and dissolving the polymer, a photoacid generator (designated as PAG1 and 2), a dissolution regulator (designated as DRR1 to 4), a basic compound, and a compound having a $\equiv\text{C}-\text{COOH}$ group in the molecule (ACC1 and 2) in a solvent in accordance with the formulation shown in Tables 1 to 3. These compositions were each filtered through a Teflon filter (pore diameter 0.2 μm), thereby giving resist solutions.




(PAG 1)




(PAG 2)




(DRR 1)




(DRR 2)




(DRR 3)



(DRR 4)



(ACC 1)



(ACC 2)

The solvent and basic compounds used are as follows.
It is noted that the solvent contained 0.01% by weight of
surfactant KH-20 (Asahi Glass Co., Ltd.).

PGMEA: propylene glycol methyl ether acetate

TEA: triethanolamine

TMMEA: trimethoxymethoxyethylamine

TMEMEA: trimethoxyethoxymethoxyethylamine

These resist solutions were spin coated onto silicon
wafers having an antireflection film (ARC25 by Nissan
Chemical Co., Ltd., 77 nm thick) coated thereon, then heat
treated at 130°C for 90 seconds to give resist films having
a thickness of 375 nm. The resist films were exposed using
an ArF excimer laser stepper (Nikon Corporation; NA 0.55),
then heat treated at 110°C for 90 seconds, and puddle
developed with a solution of 2.38% tetramethylammonium
hydroxide in water for 60 seconds, thereby giving 1:1 line-
and-space patterns.

The wafers as developed were sectioned and observed
under sectional SEM. The optimal dose (Eop, mJ/cm²) was

defined as the dose which provided a 1:1 resolution at the top and bottom of a 0.20 μm line-and-space pattern. The resolution of the resist under evaluation was defined as the minimum line width (μm) of the lines and spaces that

- 5 separated at the optimal dose. The shape of the resolved resist pattern was examined under a SEM and classified into rectangular, rounded head, T-top, forward taper or reverse taper.

10 The composition and test results of the resist materials are shown in Tables 1 to 3.

10073223.021302

Table 1

| Example | Resin (pbw) | Photoacid generator (pbw) | Dissolution regulator (pbw) | Basic compound (pbw) | Solvent (pbw) | Eop, mJ/cm ² | Resolution, μ m | Shape |
|---------|-----------------|---------------------------|-----------------------------|----------------------|---------------|-------------------------|---------------------|-----------------------|
| 1 | Polymer 1 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 36.0 | 0.16 | rectangular |
| 2 | Polymer 2 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 38.0 | 0.16 | rectangular |
| 3 | Polymer 3 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 34.0 | 0.16 | rectangular |
| 4 | Polymer 4 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 36.0 | 0.16 | rectangular |
| 5 | Polymer 5 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 36.0 | 0.16 | rectangular |
| 6 | Polymer 6 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 34.0 | 0.16 | rectangular |
| 7 | Polymer 7 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 38.0 | 0.16 | rectangular |
| 8 | Polymer 8 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 34.0 | 0.17 | rectangular |
| 9 | Polymer 9 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 32.0 | 0.16 | rectangular |
| 10 | Polymer 10 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 30.0 | 0.16 | rectangular |
| 11 | Polymer 11 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 34.0 | 0.16 | rectangular |
| 12 | Polymer 12 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 36.0 | 0.16 | rectangular |
| 13 | Polymer 1 (80) | PAG 2 (1) | - | TEA (0.125) | PGMEA (480) | 38.0 | 0.16 | rectangular |
| 14 | Polymer 1 (80) | PAG 2 (1) | - | TMMEA (0.236) | PGMEA (480) | 38.0 | 0.15 | rectangular |
| 15 | Polymer 1 (80) | PAG 2 (1) | - | TMEMEA (0.347) | PGMEA (480) | 40.0 | 0.16 | rectangular |
| 16 | Polymer 5 (70) | PAG 2 (1) | DRR 1 (10) | TMMEA (0.236) | PGMEA (480) | 32.0 | 0.17 | somewhat rounded head |
| 17 | Polymer 5 (70) | PAG 2 (1) | DRR 2 (10) | TMMEA (0.236) | PGMEA (480) | 34.0 | 0.16 | rectangular |
| 18 | Polymer 5 (70) | PAG 2 (1) | DRR 3 (10) | TMMEA (0.236) | PGMEA (480) | 40.0 | 0.16 | rectangular |
| 19 | Polymer 5 (70) | PAG 2 (1) | DRR 4 (10) | TMMEA (0.236) | PGMEA (480) | 34.0 | 0.16 | rectangular |
| 20 | Polymer 6 (80) | PAG 2 (1) | ACC 1 (4) | TMMEA (0.236) | PGMEA (480) | 32.0 | 0.17 | somewhat rounded head |
| 21 | Polymer 6 (80) | PAG 2 (1) | ACC 2 (4) | TMMEA (0.236) | PGMEA (480) | 36.0 | 0.16 | rectangular |

Table 2

| Example | Resin (pbw) | Photoacid generator (pbw) | Dissolution regulator (pbw) | Basic compound (pbw) | Solvent (pbw) | Eop, mJ/cm ² | Resolution, μ m | Shape |
|---------|-----------------|---------------------------|-----------------------------|----------------------|---------------|-------------------------|---------------------|-----------------------|
| 22 | Polymer 13 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 36.0 | 0.16 | rectangular |
| 23 | Polymer 14 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 38.0 | 0.16 | rectangular |
| 24 | Polymer 15 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 34.0 | 0.16 | rectangular |
| 25 | Polymer 16 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 32.0 | 0.16 | rectangular |
| 26 | Polymer 17 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 32.0 | 0.16 | rectangular |
| 27 | Polymer 18 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 28.0 | 0.16 | rectangular |
| 28 | Polymer 19 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 36.0 | 0.15 | rectangular |
| 29 | Polymer 20 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 38.0 | 0.16 | rectangular |
| 30 | Polymer 13 (80) | PAG 2 (1) | - | TEA (0.125) | PGMEA (480) | 38.0 | 0.16 | rectangular |
| 31 | Polymer 13 (80) | PAG 2 (1) | - | TMMEA (0.236) | PGMEA (480) | 38.0 | 0.15 | rectangular |
| 32 | Polymer 13 (80) | PAG 2 (1) | - | TMMEA (0.347) | PGMEA (480) | 40.0 | 0.16 | rectangular |
| 33 | Polymer 14 (70) | PAG 2 (1) | DRR 1 (10) | TMMEA (0.236) | PGMEA (480) | 36.0 | 0.17 | somewhat rounded head |
| 34 | Polymer 14 (70) | PAG 2 (1) | DRR 2 (10) | TMMEA (0.236) | PGMEA (480) | 38.0 | 0.16 | rectangular |
| 35 | Polymer 14 (70) | PAG 2 (1) | DRR 3 (10) | TMMEA (0.236) | PGMEA (480) | 44.0 | 0.16 | rectangular |
| 36 | Polymer 14 (70) | PAG 2 (1) | DRR 4 (10) | TMMEA (0.236) | PGMEA (480) | 38.0 | 0.16 | rectangular |
| 37 | Polymer 15 (80) | PAG 2 (1) | ACC 1 (4) | TMMEA (0.236) | PGMEA (480) | 34.0 | 0.16 | somewhat rounded head |
| 38 | Polymer 15 (80) | PAG 2 (1) | ACC 2 (4) | TMMEA (0.236) | PGMEA (480) | 38.0 | 0.16 | rectangular |

Table 3

| Example | Resin (pbw) | Photoacid generator (pbw) | Dissolution regulator (pbw) | Basic compound (pbw) | Solvent (pbw) | Eop, mJ/cm ² | Resolution, μm | Shape |
|---------|--------------------|---------------------------------|-----------------------------------|----------------------------|------------------|----------------------------|-------------------|--------------------------|
| 39 | Polymer 21 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 34.0 | 0.16 | rectangular |
| 40 | Polymer 22 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 36.0 | 0.16 | rectangular |
| 41 | Polymer 23 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 36.0 | 0.16 | rectangular |
| 42 | Polymer 24 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 38.0 | 0.16 | rectangular |
| 43 | Polymer 25 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 32.0 | 0.16 | rectangular |
| 44 | Polymer 26 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 30.0 | 0.16 | rectangular |
| 45 | Polymer 27 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 34.0 | 0.16 | rectangular |
| 46 | Polymer 28 (80) | PAG 1 (1) | - | TEA (0.125) | PGMEA (480) | 36.0 | 0.16 | rectangular |
| 47 | Polymer 21 (80) | PAG 2 (1) | - | TEA (0.125) | PGMEA (480) | 36.0 | 0.16 | rectangular |
| 48 | Polymer 21 (80) | PAG 2 (1) | - | TMMEA (0.236) | PGMEA (480) | 36.0 | 0.15 | rectangular |
| 49 | Polymer 21 (80) | PAG 2 (1) | - | TMEMEA (0.347) | PGMEA (480) | 40.0 | 0.15 | rectangular |
| 50 | Polymer 22 (70) | PAG 2 (1) | DRR 1 (10) | TMMEA (0.236) | PGMEA (480) | 34.0 | 0.17 | somewhat rounded head |
| 51 | Polymer 22 (70) | PAG 2 (1) | DRR 2 (10) | TMMEA (0.236) | PGMEA (480) | 36.0 | 0.16 | rectangular |
| 52 | Polymer 22 (70) | PAG 2 (1) | DRR 3 (10) | TMMEA (0.236) | PGMEA (480) | 40.0 | 0.16 | rectangular |
| 53 | Polymer 22 (70) | PAG 2 (1) | DRR 4 (10) | TMMEA (0.236) | PGMEA (480) | 36.0 | 0.16 | rectangular |
| 54 | Polymer 23 (80) | PAG 2 (1) | ACC 1 (4) | TMMEA (0.236) | PGMEA (480) | 32.0 | 0.17 | somewhat rounded head |
| 55 | Polymer 23 (80) | PAG 2 (1) | ACC 2 (4) | TMMEA (0.236) | PGMEA (480) | 36.0 | 0.16 | rectangular |

It is seen from Tables 1 to 3 that the resist
5 compositions within the scope of the invention have a high
sensitivity and resolution upon ArF excimer laser exposure.

Japanese Patent Application Nos. 2001-037247, 2001-
037262 and 2001-037271 are incorporated herein by reference.

Although some preferred embodiments have been described, many modifications and variations may be made thereto in light of the above teachings. It is therefore to be understood that the invention may be practiced otherwise
5 than as specifically described without departing from the scope of the appended claims.

10073223.021302